

**Teratogenic Properties of 4(3H) Quinazolinone-2-Propyl-2-Phenyl Ethyl on Stomach and Heart Development of New Born Balb/C Mice**

**Nasim Javdan, Jasem Estakhr**

**Science and Research Branch, Islamic Azad University, Fars, Iran.**

\* **Corresponding Address:** Jasem Estakhr, [j.estakhr@yahoo.com](mailto:j.estakhr@yahoo.com). Tel: +989179283966.

**Summary**

Quinazolinones are used for treatments of prevalent diseases. They belong to hyponic and potent anticonvulsant drugs, Act strongly, inhibiting human immunodeficiency virus (HIV) and cancer, Enter circulatory system and pass through placenta barrier. In this study, for the first time different aspects of developmental effects of 4(3H) quinazolinone-2- propyl-2-phenyl ethyl (QPPE) on stomach and heart of Balb/C mice embryos were investigated. Pregnant Balb/C mice were divided into 3 groups (n= 10) of control, receiving distilled water, sham, receiving 0.05% methyl cellulose (the solvent) and experimental group, receiving one of the most effective dose of 100 mg/kg/body weight of QPPE, by IP injections on day 8th to 15th of gestation. After anesthetising mothers, stomachs and hearts of 5-day old newborn Balb/C mice were removed, fixed and stained with H & E for light microscopic and quatitative studies. Results showed symptoms of gastritis (hyperaemia and decrease in thickness of mucus layer) in newborn Balb/C mice of treated groups. QPPE also created necrotic cells and an increase in connective tissues of hearts of newborn mice of treated groups. In conclusion by being teratogens and toxins, these two new derivatives affected development of embryonic stomach and heart at histological level.

**Introduction**

Different teratogens and toxins have different developmental effects on different species with different severity (1, 2). By passing through placenta (3, 4), treatments with teratogens such as alcohol, quinazolinones, etc create early death, malformations and irregularities in different parts of developing embryos. Quinazolinones are heterocyclic components with various characteristics; such as: anti-inflammation, anti-malaria, anti-spasm, anti-microbial, anti-hypertensive, anti-allergic, sedative, anti-tuberculosis, antihyperlipidemic, anxiolytic, analgesic, anticonvulsant, as well as hypnotic activities. They are also known for their fungicidal properties, inhibition of tyrosine-kinase (involved in tubulin and 8-hydroxy-2-methyl quinazolinone polymerization), DNA repair enzyme poly (ADP-ribose) polymerase (PARP), and hhs signalling pathways. They are also effective in treatment of osteoarthritis, cancer, diabetes and parkinsonism complications (5-8). Previous studies showed that 4(3H)-quinazolinone-2-propyl-2-phenylethyl (QPPE) can causes morphological, skeletal and histological abnormalities in Balb/C mice embryos (9-14). The mechanisms of the effects of quinazolinones on embryonic cells are not clear yet, but there are few reports showing its toxic characteristics. Following our earlier demonstrations of their toxic effects at morphological and skeletal levels of Balb/C mice fetuses and embryos (15, 16), the pathological effects of QPPE have been investigated on the morphological and histological structures of newborn Balb/C mice internal organs, such as stomach, heart.

### Materials and methods


Balb/C mice (8 to 12 weeks old) were purchased from Razi Institute,(Karaj, Iran), weighing 27-28 g were used in this study. Animals were maintained under a 12:12-hour light/dark photoperiod. Female mice were mated with males of the same strain (1:2) and isolated the following morning, upon finding the vaginal plug, day zero of the pregnancy was designated and mated animals were kept singly in cages, at ambient room temperature.


The new derivative of quinazolinones: 4(3H)-quinazolinone-2-propyl-2-phenylethyl (QPPE), synthesized at Department of Chemistry , Faculty of Science , University of Shahid-Beheshti , Tehran , Iran (17) were used for IP injection. So, pregnant mice were divided into 3 groups (n=10) of control, sham, and experimental, received distilled water (10ml/kg), methyl cellulose %0.05 (10ml/kg) (the solvent of quinazolinones) and 100 mg/kg Balb/C body weight of QPPE (most effective dose), respectively, by IP injection, on days 8th to 15th of gestation. 5day old newborns were killed by cervical dislocation. Hearts and stomachs of 4-days old Balb/C mice were removed, fixed in %10 formaldehyde, stained with H and E, studied with light microscope. Parametric data were analyzed by statistical packages for social sciences (SPSS, version 9.0).

### Results


No morphological abnormalities were observed in the stomachs of newborn Balb/C mice of experimental group in comparison with sham and control groups. There were no significant differences between control and sham groups. Injection of 100mg/kg/body weights of QPPE, resulted in the formation of normal (without hyperaemia) and abnormal (with hyperaemia) stomachs in newborn Balb/C mice. As statistical analysis revealed, significant decrease occurred in the thickness of mucosal layer of stomachs of newborn Balb/C mice of experimental group 1 (Figs. 1 & 2), comparing with stomachs of newborn Balb/C mice of sham and control groups. There were no significant differences between mucosal layers of stomachs of newborn Balb/C mice of QPPE group. No significant differences were observed between morphological and histological structures of hearts of newborn Balb/C mice of control and sham groups. Injection of 100mg/kg/body weights of QPPE created abnormal hearts, hearts with necrotic cells and connective tissues between myocytes of newborn Balb/C mice comparing with newborn Balb/C mice hearts of sham and control groups (Figs. 3 & 4).

**Figure 1.** Comparison of the thickness (Pm)of mucus layer of stomachs of four different groups of newborn Balb/C mice. QEPE was more effective ( $P<0.05$ ).






**Figure 2.** Hyperaemia(white arrows) and decrease in the thickness of mucus layers(black arrows) of stomachs of groups treated with QPPE(40X).



**Figure 3.** Necrosis (arrows) in hearts of newborn Balb/C mice of mother treated with QPPE.



**Figure 4.** Connective tissues(arrows) in hearts of newborn Balb/C mice of mother treated with QPPE.

### **Discussion**

Results of some pharmacokinetic studies suggested that quinazolinones pass through placenta barrier by simple diffusion, along a chemical gradient. Two possible explanations for this passage were considered: first possibility is related to some chemicals directly interacting with endogenous receptors for hormones, growth factors, cell signalling molecules and other endogenous compounds. Second possibility shows that receptors can be broadly classified as cytosolic/nuclear or membrane bound. As results demonstrated, quinazolinones pass through placenta barrier by simple diffusion, along a chemical gradient entering gastric tissues, causing inflammation (gastritis) , because of atrophy in mucus layer, likely as the result of creating necrosis in stomach cells. QPPE and QEPE are without active chemical groups but generate free radicals and active metabolites after metabolization leading to lipid peroxidation ,destroying cell membranes after releasing intracellular components such as lysosomal enzymes, causing further tissue damages. Oxygen derived free radicals play pathological roles in gastritis and radical scavengers such as alfa tocopherol, carotenoids glutathione redox system play a significant role in protecting membranes from oxidative damages. Depletion of gastric mucus GSH may result in accumulation of free radicals, initiating membrane damages by lipid peroxidation, ultimately leading to necrosis in specially parietal cells ,because of their numerous surface receptors and pumps specializing for gastric acid production(18). Apoptosis is uncommon in normal mucosal layer ,but chronic inflammation is associated with increased apoptosis, happening mainly and only in mucus surface (19) . Appearance of connective tissues between myocytes is due to the stimulation of the multiplication of fibroblasts because of necrosis and their annihilation,so, necrotic heart cells will be replaced by connective tissue (20).

### **References**

1. Derelanko MJ, Hollinger MA. Handbook of toxicology. Boca Raton: CRC Press; 2002.
2. Gardella JR, Hill JA. Environmental toxins associated with recurrent pregnancy loss. *Semin Repro Med* 2000; 18:407-424.
3. Fowden AL, Ward JW, Wooding FPB, Forhead AJ, Constancia M. Programming placental nutrient transport capacity. *J Physiol* 2006; 572:5-15.

4. Coan PM, Angiolini E, Sandovici I, Burton GJ, Constancia M, Fowden AL. Adaptations in placental nutrient transfer capacity to meet fetal growth demands depend on placental size in mice. *J Physiol* 2008; 15:4567-4576.
5. Abdel-Alim M, El-Shorbag Abdel-Nasser A, El-Gendy Mahmoud A, El-Shareif Hosny AH. Quinazolinone derivatives of biological interest: V. Novel 4(3H)-Quinazolinones with sedative-hypnotic, anticonvulsant and anti-inflammatory activities. *Coll Cze Che Comm*2008; 58:1963-1968.
6. Jiang S, Zeng Q, Gettayacamin M, Tungtaeng A, Wannaying S, Lim A, *et al.* Anti-malaria activities and therapeutic properties of febrifugine analogs. *Antimicrob Agents Chemother* 2005; 49:1169-1176.
7. Refaie FM, Esmat AY, Abdel Gawad SM, Ibrahim AM, Mohamed MA. The anti-hyperlipidemic activities of 4(3H) quinazolinone and two halogenated derivatives in rats. *Lipids Health Dis* 2005; 4:22-30.
8. Yadav MR, Shirude ST, Parmar A, Balaraman R, Giridhar R. Synthesis and anti-inflammatory activity of 2,3- diaryl-4(3H)-quinazolinones. *Chem Heter Com* 2006; 42:1038-1045.
9. Shams lahijani M, Ahmadzadeh F, Dabiri M. Teratogenic effects of new quinazolinone derivative on the development of Balb/C mice fetuses on days 9, 10 and 11 of gestation. *Journal of Science and Technology* 2006; 30A1: 1-8.
10. Shams lahijani M, Aounegh R. Teratogenic effect of quinazolinone on Balb/C mice fetuses. *Journal of Medical Sciences Research* 2007; 1 (1).
11. Etemad S, Shams Lahijani M. Quinazolinones and nephrotoxicity in new born Balb/C mice. 7th World Congress of Nephrology(WCN) Rio De Janeiro Brazil 2007.
12. Fadavi M, Shams Lahijani M. Pathological effects of quinazolinones on the small intestine of new born Balb/C mouse. XI International Congress of Toxicology (ICT) Montreal Canada 2007.
13. Rajabi H, Shams Lahijani M. Histological study of liver of newborn Balb/C mice treated with quinazolinones. XI International Congress of Toxicology (ICT) Montreal Canada 2007.
14. Estakhr Jasem, Sanchooli Naser, Hatami Leili, Shams Lahijani Maryam. Investigation of the immunopathologic effects of 4(3H)-quinazolinone-2-propyl-2-phenyl ethyl (QPPE) in new born Balb/C mice. *Pharmacologyonline*, 2009; 1: 377-384.
15. Shams Lahijani M., Ahmadzadeh F., Dabiri M. Teratogenic effects of a new derivative of quinazolinone on the development of Balb/C mice embryos, on days 9, 10 and 11 of gestation. *Ir. J. Sci. Technol.* 2006;30: 1- 8.
16. Shams Lahijani M., Aounagh R. Teratogenic effects of quinazolinones on Balb/C mice fetuses. *J. Med. Sci. Res.* 2007;1,25-30.
17. Dabiri M, Salehi P, Khajavi MS, Mohammadi A. Microwave-assisted one-pot three component synthesis of some new 4(3H)-quinazolinone derivatives. *Heterocycles* 2004; 63(6):1417-21.
18. Bayiroğlu F., Cemek M., Caksen H., Cemek F., Dede S. Altered antioxidant status and increased lipid peroxidation in children with acute gastroenteritis admitted to a pediatric emergency service. *J. Em. Med.* 2009; 36( 3):227-31.
19. Van Grieken N.C.T., Meijer G.A., Hausen A., Meuwissen S.G.M., Baak J.P.A., Kuipers E.J. Increased apoptosis in gastric mucosa adjacent to intestinal metaplasia. *J. Clin. Pathol.* 2003; 56(5): 358–61.
20. Cohen-Gould L., Robinson T.F., Factor S.M. Intrinsic connective tissue abnormalities in the heart muscle of cardiomyopathic Syrian hamsters. *Am. J. Pathol.* 1987; 127(2): 327–34.