

**Anti-Diarrheal Potential of *Aegle Marmelos* Corr. Root Bark Extract in Rats**

Vrushabendra Swamy. B.M \*, Nataraj KS<sup>1</sup>, Jayaveera K.N<sup>2</sup>, Gopkumar P<sup>3</sup>. Sridevi Nayak<sup>3</sup>, Kumar G.S<sup>4</sup>, Sanjay P Umachigi.<sup>4</sup>

\* Rural College of Pharmacy, Devanahalli-562 110 Karnataka, India.

<sup>1</sup> SRR College of Pharmaceutical Sciences, Valbhapur Village, Elkhathurthy Mandal, Karim Nagar Dist - 505476, Andhra Pradesh, India.

<sup>2</sup>JNTU, Ananthpur 515 001, Andhra Pradesh, India.

<sup>3</sup>Department of Chemical Engineering, National Institute of Technology Karnataka , Surathkal 575 025, Karnataka, India.

<sup>4</sup> Rural College of Pharmacy, Devanahalli-562110 Karnataka, India.

**Summary**

A study was undertaken to evaluate the effect of methanolic extract of the root-bark of *Aegle marmelos* Corr. (MAMC) against several experimental models of diarrhoea in rats. MAMC treated animal's showed significant inhibitory effect against castor-oil induced diarrhea and PGE<sub>2</sub> induced enteropooling in rats. The extract also showed a significant reduction in gastrointestinal motility in the charcoal meal test in rats. The results obtained to establish the efficacy and substantiate the folkare claim as an anti-diarrhoeal agent.

Key words: *Aegle marmelos*, MAMC, root-bark, diarrhea, methanol extract.

**\*For correspondence:**

Vrushabendra Swamy B.M  
Department of Pharmacology,  
Rural College of Pharmacy,  
P.B No 10, D.S.Road,  
Devanahalli, Bangalore,  
Karnataka, India. -562110  
Phone: 0091 80 7682450  
Fax: 0091 80 7682450  
**E-mail:** swamybm@yahoo.com

**Introduction**

Diarrhoea has long been recognized as one of the most important health problem in the developing countries (1). World Wide distribution of diarrhea accounts for more

than 5-8 millions deaths each year in infants and small children's less than 5 years. According to WHO estimation for the year 1998, there were about 7.1 million deaths due to diarrhea (2). Secretory diarrhoea is most dangerous symptom of gastrointestinal problems (3) and is associated with excessive defecation and stool outputs. The stool being of abnormally loose consistency (4). The World Health Organization has constituted a Diarrhoeal Disease Control program (CDD), which includes studies of traditional medicinal practices, together with the evaluation of health educational and prevention approaches (5).

*Aegle marmelos* Corr. (Rutacea), known as Bael in Hindi, Sripal or Bilwa in Sanskrit and Bael tree in English (6). Traditionally this plant useful in treating pain, fever, inflammation, respiratory disorders, cardiac disorders, dysentery and diarrhea (7-9). Reports indicate the pharmacological activities of this plant include anti-bacterial & antifungal (10-12), hypoglycemic (13), anticancer (14), cardio tonic (15) and anti-inflammatory, anti-pyretic and analgesic activity (16). The phytochemical constituents of root-bark contain tannins, reducing sugars (6), alkaloids, sterols, coumarins and essential oils (17).

## **Materials and Methods**

### **Plant Material**

The root bark of *Aegle marmelos* Corr. were collected from Nandhi hills, Kolar district, Karnataka, India in February 2005. The plant was identified by routine pharmacognostical studies including organoleptic tests, microscopic and macroscopic observation. The vouches specimen AMC1/ RCP 04-05 has been retain in our laboratory for future reference. The collected root barks were air dried and pulverized using mechanical grinder.

### **Preparation of Extract**

The powdered root-barks (500g) were extracted using 85% methanol. The solvent was removed completely removed from the extract by distillation *in vacuo* and a brown

coloured semisolid mass obtained. (Yield 16.2% w/w, with respect to the dry powder). The extract was stored in dessicator and used for further experiment after suspend in aqueous tween 80 solution (2%). The chemical constituents of the extracts were identified by qualitative tests and further confirmed by TLC study for the presence of alkaloids, tannins, sterols, and carbohydrates.

### **Animals**

Wister Albino rats (either sex) weighing between 150-200gm were used. The animals were divided into five groups (n=6) and were housed in polyacrylic cages with standard diet and water *ad libitum* except in the castor oil model, in which the animals were placed individually in separate cages. The animals were maintained under standard laboratory conditions for acclimatization period of 7 days prior to perform the experiment. The animals were fasted for 18hours prior to the experiment.

## **Experimental Design**

### **Toxicity Study**

An acute toxicity study relating to the determination of LD<sub>50</sub> value was performed using different doses of the extract according to the method described by Ghoshet.al (18) from the toxicity study, it was observed that the root-bark extract is non-toxic upto dose of 3.2g/kg body weight. It is safe and was used in different doses for further studies (Table 2).

### **Castor Oil Induced Diarrhoea (19)**

The doses of MAMC were selected on a trial basis and administered orally (100, 200 & 400mg/kg body weight) by gavage to three groups of animals. The fourth group received diphenoxylate (5mg/kg body weight) orally and the fifth group received neither drug nor extract but 2% v/v aqueous Tween 80 (1 ml) only and served as a control. After 60 min of drug treatment, each animal was administered 1ml of castor oil orally by

gavage and observed for defecation up to 4hrs after castor oil administration. Characteristic diarrheal droppings were noted in the transparent plastic dishes placed beneath the individual perforated rat cages. The mean number of wet feces was calculated from the diarrhoeal droppings in the transplant plastic dishes (20).

#### **PGE<sub>2</sub> – Induced Enteropooling (21)**

For this evaluation, rats of same stock as above were deprived of food and water for 18hrs prior to the experiment. Five groups of six animals were used, which were placed in five perforated cages. The first three groups of rats were treated with MAMC (100, 200 & 400mg/kg body weight, p.o) while the fourth and fifth group received 1ml of 5% v/v ethanol in normal saline (i.p). The fourth group was then administered 1ml of normal saline and used as control. Immediately afterwards, each rat was treated with PGE<sub>2</sub> (100µg/kg body weight in 5% v/v ethanol in normal saline) administered orally. All the rats were sacrificed under mild anesthesia after 30min. The entire length of intestine from the pylorus to the caecum was dissected out, and its contents were collected and measured.

#### **Gastrointestinal Motility Test (21)**

In this method rats were fasted for 18hrs and placed in five metal cages, six in each. Each animal was given 1ml of charcoal meal (3% deactivated charcoal in normal saline). The first three groups of animals were administered MAMC orally (100, 200 & 400 mg/kg body weight) immediately after the charcoal meal treatment. The fourth group received atropine (0.1mg/kg body weight, i.p) as standard for comparison. The fifth group was treated with normal saline as control. 30min after administration of the charcoal meal, animals of each individual group were killed and the movement of charcoal from pylorus to caecum was measured. The charcoal movement in the intestine was expressed as percentage.

#### **Statistical Analysis**

For all the above experiments results were expressed as mean  $\pm$ SEM. Statistical significance tests were performed using the students't' test and p-values (Graph Pad software) were calculated by comparison with control groups.

## Results

### Chemical Analysis

The results of the preliminary photochemical screening of methanolic extract of *Aegle marmelos* corresponding root-bark have been presented in Table – 1.

**Table-1:** Phytochemical screening of *Aegle marmelos* Corr, -, absence; +, presence.

Test	Methanol extract
Alkaloids	+
Tannins	+
sterols	+
Carbohydrates	+

### Toxicity Study

From the toxicity study it was observed that the root-bark extract is non-toxic and caused no death up to a dose of 3.2g/kg orally. It is safe and was used in doses for further studies (Table – 2)

**Table-2: Toxicity Study:**

Treatment	Dose (mg/kg body weight)	No. of animals	No. of survives	No. of deaths	LD <sub>50</sub> valve
Control	2% tween 80 solution	20	20	0	-
MAMC	100	20	20	0	-
	200	20	20	0	-
	400	20	20	0	-
	800	20	20	0	-
	1600	20	20	0	-
	3200	20	20	0	>3.2g/kg body weight

MAMC: Methonolic extract of *Aegle marmelos* corr.

**Inhibition of Castor-Oil Induced Diarrohea**

The extract (MAMC) inhibited the frequency of defecation significantly, like standard drug (diphenoxylate) as compared to control (2% aqueous tween 80 treated). The wetness of fecal material also reduced by both the standard and extract (MAMC). The results are shown in Table – 3.

**Table-3:** Inhibition of castor oil induced diarrhea.

<b>Oral pretreatment at 60min (mg/kg body weight)</b>	<b>Mean number of wet feces (mean ± SEM)</b>
2% v/v aqueous tween 80(5)	27.2 ± 2.01
Diphenoxylate(5)	10.8 ± 1.44 <sup>a</sup>
MAMC(100)	21.3± 1.63 <sup>b</sup>
MAMC(200)	15.6 ± 1.17 <sup>a</sup>
MAMC(400)	12.8 ± 1.09 <sup>a</sup>

<sup>a</sup> P<0.001, <sup>b</sup> p<0.01 as compared to control (n=6)

MAMC: Methanol extract of *Aegle marmelos* Corr.

**Anti-Enteropooling Activity**

PGE<sub>2</sub> induced a significant increase in the fluid volume of the rat as compared to control animals receiving only ethanol in normal saline. The extract (MAMC) significantly inhibited PGE<sub>2</sub> induced enteropooling in rats at almost all doses used (Table – 4).

**Table-4:** Anti-enteropooling activity

Treatment	Volume of intestinal fluid ml(mean $\pm$ SEM)	P – value
Ethanol in saline	0.76 $\pm$ 0.13	-
PGE2 in ethanol	2.93 $\pm$ 0.19	<0.001 <sup>a</sup>
MAMC(100mg/kg body weight)	1.84 $\pm$ 0.11	<0.001 <sup>b</sup>
MAMC(200mg/kg body weight)	1.22 $\pm$ 0.12	<0.001 <sup>b</sup>
MAMC(400mg/kg body weight)	1.07 $\pm$ 0.05	<0.001 <sup>b</sup>

<sup>a</sup> significance with respect to ethanol in saline treatment

<sup>b</sup> with respect to PGE<sub>2</sub> treatment (n=6)

MAMC : methanol extract of *Aegle marmelos* corr.

### Effect on Gastrointestinal Motility

The extract (MAMC) decreased propulsion of charcoal meal through the gastrointestinal tract significantly with respect to the control group. The effect was comparable to the standard drug. The results are shown in Table – 5.

**Table-5:** Effect on gastrointestinal motility

Treatment	Dose (mg/kg body weight)	Movement of charcoal meal (%)
Control	2% aqueous tween 80	85.2 $\pm$ 2.1
Atropine	0.1	40.6 $\pm$ 2.3 <sup>a</sup>
MAMC	100	74.4 $\pm$ 2.5 <sup>b</sup>
MAMC	200	65.6 $\pm$ 2.1 <sup>a</sup>
MAMC	400	54.8 $\pm$ 2.3 <sup>a</sup>

P-value calculated with respect to control group (n=6)

<sup>a</sup> p<0.001, <sup>b</sup> p<0.02

MAMC: Methanol extract of *Aegle marmelos* Corr.

### Discussion

In developing countries a quarter of infant and childhood mortality is related to the diarrhea (22). The highest mortality rates have been reported to be in children less than five years of age. During the past decade oral dehydration therapy has reduced mortality from acute diarrheal disease, whereas chronic diarrhea remains a life-threatening problem in those regions in which malnutrition is common co-existing and complication factors number of factors, such as infective, immunological and nutritional has been involved in the perpetuation of the diarrheal syndrome (23). Many plants conveniently available in India are used in traditional folklore medicine for the treatment of diarrhea and dysentery of the indigenous plants used, *Andrographis Peniculata*, *Asparagus racemosus*, *Butea monosperm*, *Cassia auriculata* and other are mentioned (24). Several studies have shown that prior administration with some plant extracts had protective effect on the intestinal tract (25,26,27).

In the present study, methanol extract of root-bark of *Aegle marmelos* have not been studied so far was evaluated for its anti-diarrhoeal potential against castor oil induced diarrhea, gastro intestinal motility in charcoal meal test & PGE<sub>2</sub> induced enteropooling in Albino Wistar rats.

The MAMC exhibited significant anti-diarrheal activity against castor oil induced diarrhea in rats. The extract had a similar activity as diphenoxylate, when tested at 100, 200 & 400mg/kg and statistically significant reduction in the frequency of defecation and the wetness of the fecal droppings when compared to untreated control rats.

It is widely known that castor oil or its active component ricinoleic acid induces permeability changes in mucosal fluid and electrolyte transport that result in hyper secretory response of diarrhea (28,29). The experimental studies in rat's demonstrated a significant increase in the portal venous PGE<sub>2</sub> concentration following oral administration of castor oil (30). The ricinoleic acid from castor oil results in irritation and inflammation of the intestinal mucosa, leading to release of prostaglandins, which stimulate motility and secretion (31). Inhibitors of prostaglandin biosynthesis delayed castor oil induced diarrhea (19).

The MAMC significantly inhibited the PGE<sub>2</sub> induced intestinal fluid accumulation (enteropooling). It has been shown that E type of prostaglandin cause

diarrhoea in experimental animals as well as human beings (32). Their mechanism has been associated with dual effects on gastrointestinal motility as well as on water and electrolyte transport (33). PGE<sub>2</sub> also inhibit the absorption of glucose a major stimulus to intestinal adsorption of water and electrolytes (34).

These observations tends to suggest that MAMC reduced diarrhea by inhibiting PGE<sub>2</sub> induced intestinal accumulation of fluid.

The MAMC appears to act on all parts of the intestine. Thus it reduced the intestinal propulsive movement in the charcoal meal treated model. The MAMC showed activity similar to that of atropine. Previous study shows that activated charcoal avidly absorbs drugs and chemicals on the surface of the charcoal particles there by preventing absorption (35). Thus gastrointestinal motility test with activated charcoal was carried out to find out the effect of MAMC on peristalsis movement. The results shows that the MAMC suppressed the propulsion of charcoal meal thereby increased the absorption of water and electrolytes.

Previous reports have demonstrated the antidiarrhoeal activity of tannins (36), alkaloids (37), sterols and reducing sugars (38) containing plant extracts. The phytochemical analysis of the extract showed presence of tannins, alkaloids sterols and reducing sugar. These constituents may responsible for the antidiarrhoeal activity.

### **References**

1. Snyder JD, Merson MH. The magnitude of the global problem of acute diarrhea disease: A review of active surveillance of data. Bull WHO 1982;60:605-613.
2. Park K. Park's text book of Preventive and Social Medicine. Banarsidas Bharat: Publishers: Jabalpur, 2000:122-175.
3. Fontaine O. Diarrhea and treatment. Lancet 1988;28:1234-1235.
4. Aranda MJ, Gianella RA. Acute diarrhea: A practical review AM J Med 1999;106:670-676.
5. Pulok K, Mukherjee J, Das R, Balasubramanian, Kakali Saha, Pal M, Saha BP. Antidiaraheal evaluation of *Nelumba nueofera* rhizome extract. Ind J Pharmcol 1995;27:262-65.

6. Nadkarni AK. Dr. KM. Nadkarni's Indian Materia Medica. Vol-1. Popular Prakashan Company: Bombay: India, 1986:45.
7. Dymock WC.J, Warden H, David H. Pharmacographia indica. In: A history of the principal drugs of vegetable Origin Met within British India. Vol-1. Kegan paul: Trench Trubner and co. Ltd: London, 1890;277-281.
8. Kirtikar KR, Basu BD. ICS. Indian medicinal plants, 2<sup>nd</sup> ed. International book distributors: Dehradun:India, 1935:448-502.
9. Murugesu MKS. Materia Medica of Tamil system of medicine. Part-1. Tamil Nadu Siddha Medical Board: Govt of Tamil Nadu: India, 1988:638-642.
10. Banerji N, Kumar R. Studies on the seed oil of *Aegle marmelos* and its effects on some bacterial species. J Inst Chem. 1980;52: 59-60.
11. Pattnaik S, Subramanyam VR, Kale C. Antibacterial and antifungal activity of ten essential oils *In vitro*. Microbios. 1996;86:237-46.
12. Rana BK, Singh UP, Taneja V. Antifungal activity and kinetics of inhibition by essential oil isolated from leaves of *Aegle marmelos*. J Ethnopharmacol 1997;57: 29-34.
13. Ponnachan PTC, Paulose CS, Panikkar KR. Effect of leaf extract of *Aegle marmelos* in diabetic rats. Ind J Exp Biol 1993;31:345-47.
14. Singh RP, Banerjee S, Rao AR. Effect of *Aegle marmelos* on biotransformation enzyme system and protection against free radical mediated damage in mice. J Pharm Pharmacol 2000;52: 991-1000.
15. Haravey SK. A preliminary commutation of the action of *Aegle marmelos* (Bael) on heart. Ind J Med Res 1968;56:327-331.
16. Veerappan A, Shigeru MR, Dhananjayan A. Studies on the anti-inflammatory, antipyretic and analgesic properties of the leaves of *Aegle marmelos* corr. J Ethnopharmacol 2005; 96:159-163
17. Basu D, Sen R, Alkaloids and coumarins from root bark of *Aegle marmelos*. Phytochem 1974;13:2329-30
18. Ghosh MN. Fundamental of Experimental Pharmacology. 2<sup>nd</sup> ed. Scientific book agency: Calcutta: India, 1984: 53.

19. Awouters F, Nimegeers CJE, Lenaerts FM, Janseen PAJ. Delay of castor oil diarrhea in rats; a new way to evaluate inhibitors of PG synthesis. *J Pharm Pharmacol.* 1978;30:41-45.
20. Mandal SC, Mukherjee PK, Saha K, Pal M, Saha BP. Antidiarrheal evaluation of *Ficus racemosa* Linn. Leaf extract. *J. Nat Prod Scn* 1997;3(2):100-103.
21. Gunakkunra A, Padmanahan K, Thirumal P, et al. Antidiarrheal activity of *Butea monosperma* in experimental animals. *J Ethanopharmacol* 2005;98(3):241-244.
22. Jousilahti P, Madkour SM, Lambrechts T, Sherwin E. Diarrheal disease morbidity and home treatment practical in Egypt public health, 1997;111(1):5-10.
23. Galvez J, Sanchez De Medina F, Jimenez J, et al. A effect of quercitrin on lactose- induced chronic diarrhea in rats. *Plant Med* 1995;61:302-306.
24. Chopra R.N, Nayar S.L and Chopra I.C. Glossory of Indian medicinal plants. Countil of scientific and Industrial research, New Delhi, 1956.
25. Rani S, Ahamed N, Rajaram S, et al. Antidiarrheal evaluation of clerodendrum phlomidis Linn. Leaf extract in rats. *J. ethanopharmacol* 1999; 68: 315-319,
26. Mujumdar AM, Upadhye AS, Misar AV. Studies on antidiarrhoeal activity of *Jatropha curcus* root extract in albino mice. *J Ethanopharmacol* 2000;70:183-87.
27. Kumar S, Dewan S, Sangraula H, Kumar VL. Antidiarrhoeal activity of the latex of *Calotropis procera*. *J Ethanopharmacol.* 2001;76:115-18.
28. Ammon HV, Thomas PJ, Phillips S. Effect of oleic and recinoleic acid on net jejunal water and electrolyte movement. *J Clin Inves* 1974;53:374-379.
29. Gaginella TS, Stewart JJ, Ison WA, Base P. Actions of recinoleic acid and structurally related fatty acid on gastrointestinal tract II. Effect on H<sub>2</sub>O and electrolyte absorption *In Vitro*. *J Pharmacol Expt Ther* 1975;195:355-361.
30. Luderer JR, Dermers IM, Hayes AT. Advances in prostaglandin and thromboxane research: Raven press: New York, 1980;1633-1638.
31. Peirce NF, Carpenter CCJ, Ellioh HZ, Geenough WB. Effect of transmucosal water and electrolyte movement in canine jejunum. *Gastroenterol* 1971;60:22-23.
32. Eakins KE, Sanner JM. Prostaglandins Antagonists, in Karim SMM(Ed), prostaglandins progress in research: Wiley Interscience: New York, 1972;263-64.

33. Dajani EZ, Roge EAN, Bertermann RE. Effect of E Prostaglandins, diphenoxylate and morphine on intestinal motility *In Vivo*. Eur J Pharm 1975;34:105-13.
34. Jagge BM. Prostaglandins and serotonin. Non peptide diarrhoeagenic hormones. World J Surg 1979;3:565-78..
35. Leves G. Gastrointestinal clearance of drugs with activated charcoal. New Eng J Med 1982;307:676-78.
36. Mukherjee PK. Saha K. Murugesan T, et al.. Screening of antidiarrhoeal profile of some plant extracts of a specific regions of west Bengal, India. J Ethanopharmacol 1998;60:85-89.
37. Gricilda SF, Molly T. Study of antidiarrhoeal activity of four medicinal plants in castor oil induced diarrhea. J Ethanopharmacol 2001;76: 73-76.
38. Otshudi AL, Vercruysse A, Foriers A. Contribution to the ethanobotanical, phytochemical and pharmacological studies of traditionally used medicinal plants in the treatment of dysentery and diarrhoea in Lomela area (DRC). J Ethanopharmacol 2000; 71: 411-423