

**ANTIINFLAMMATION EFFECT OF *PROSOPIS FRACTA* POD
AQUATIC EXTRACT ON NEUROGLIA DENSITY AFTER
SCIATIC NERVE INJURY IN RATS**

Maryam Tehranipour* and Mahtab Mollashahi

Department of Biology, Faculty of Science, Islamic Azad University – Mashhad
Branch, Mashhad, Iran

Summary

Neuroglial cells undergo prompt responses to peripheral motor and sensory axon injury. They participate in the pathogenesis of neurological disorders by secreting various inflammatory molecules such as cytokines or nitric oxide. One of the problems in central nervous system defeats is the return of these defeats to cellular body of those neurons which their axons are destroyed. These effects arrive to the point of defeats as retrograde and cause central degeneration in spinal cord. *Prosopis farcta* has antioxidant and antidiabetic effect. The aim of this study is carried out to examine the effects of aquatic extract of *Prosopis farcta* pod on neuroglia density after sciatic nerve injury in rats.

Thirty two adult male wistar rats were used and divided to four groups (control, compression and two experimental groups). In compression and experimental groups right sciatic nerve were highly compressed for 60 s, assigned to experimental groups (Compression + aquatic extract of *Prosopis farcta* injections (50, 75 mg kg⁻¹, ip, 2 time) (N=8).

After 4 weeks post-operative the lumbar segments of spinal cord were sampled, processed, sectioned serially and stained with toluidine blue (pH 4.65). By using stereological quantitative technique, the number of neuroglia in the right horn of spinal cord were counted and compared with each other. Statistical analyses showed remarkable increase in the number of neuroglia in the compression and all experimental groups (p<0.05). It shows that administration of *prosopis farcta* pod aquatic extract after sciatic nerve injury cannot decrease neuroglia density in compare with compression groups.

Key words: prosopis farcta, neuroglia, density, sciatic nerve

Running title: prosopis farcta aquatic extract on neuroglia density

Corresponding Author: **Tehranipour Maryam, Department of Biology, Faculty of Science, Islamic Azad University, Mashhad Branch, Mashhad, Iran. Tel: +98511835050 Fax: +985118424020E mail: maryam_tehranipour@mshdiau.ac.ir**

Introduction

Microglia are one of the three types of glial cells in the central nervous system (CNS). In contrast to neurons, astrocytes, and oligodendrocytes, they are not of ectodermal but rather of mesodermal origin. Therefore, they are closely related to other myeloid cells such as peripheral monocytes and macrophages, and are considered to be the resident macrophage-like population of the CNS. In the healthy adult, CNS microglial cells are found as so-called resting microglia, characterized by a small cell body with fine, ramified processes and low expression of surface antigens. Virtually all CNS injuries trigger a rapid activation of microglial cells (1). Such activated microglia apparently may serve as neurosupportive cells, providing neuroprotection and scavenging cellular debris in response to the injury (2).

Injured nerves regenerate slowly and often over long distances (3). Prolonged periods for regenerating nerves to make functional connections with denervated targets prolong the period of isolation of the neurons from the target (chronic axotomy) and of the denervation of Schwann cells in the distal nerve pathways (chronic denervation)(3). In an animal model, we demonstrated that prolonged axotomy and chronic denervation severely reduce the regenerative capacity of neurons to less to 10% (4). Concurrent reduction in neurotrophic factors, including brain- and glial-derived neurotrophic factors (BDNF and GDNF) in axotomized neurons and denervated Schwann cells, suggest that these factors are required to sustain nerve regeneration(5). Findings that exogenous BDNF and GDNF did not increase numbers of neurons that regenerate their axons in freshly cut and repaired rat nerves, but did increase the numbers significantly after chronic axotomy, are consistent with the view that there is sufficient endogenous neurotrophic factor supply in axotomized motoneurons and denervated Schwann cells to support nerve regeneration but that the reduced supply must be supplemented when target reinnervation is delayed (6).

Early peripheral lesions may deprive central neurons of necessary trophic factors, accentuate naturally occurring central cell death, and thereby result in smaller central representations. Or, smaller central representations may be produced by competitive interactions of deprived with adjacent intact pathways. In addition, throughout all stages of development, the capacity for reorganization may be spatially limited and depend on the size or pattern of the peripheral injury (7).

There is a long history of traditionally using the extraction of *Prosopis farcta* plant for treatment of pains in Ilam Province. Decoction of *Prosopis farcta* has also been used traditionally to reduce cardiac or chest pain in this province (8).The antioxidant activity of these plant's extracts and their potential role in radical scavenging agreed with their potential use by Jordanian population as a traditional anti-diabetic agents (9,10).

There is no report to show an academic research (*in vitro*, *in vivo*), to evaluate the protective effect of this plant so far. This study aimed to evaluate the effect of *Prosopis farcta* aquatic extract on neuroglia density after sciatic nerve injury in rat.

Material and methods

The *prosopis farcta* (herbarium code 1952) was supplied by Islamic Azad University of Mashhad, Iran (2010).

Animal subjects

Thirty two male, Wistar rats weighting between 300-350 g served as subjects for these experiments. All animals were housed individually and maintained on a 12/12 light/dark cycle, with lights on at 6.00h. Ambient temperature in the animal facility was kept at $22\pm 2^{\circ}\text{C}$. Food and water was given ad libitum.

Extraction

prosopis farcta was collected from a reign around Zahedan and was coded with Islamic Azad University of Mashhad, Iran herbarium (herbarium code 1952). For extraction 50g powder pod with 300 cc water were mixed and extraction perform with Soxhlet apparatus (11). After obtaining extract, it was situated in oven with temperature ($45^{\circ}\pm 2^{\circ}$) for 48 hours to remove solvent.

Groups

Thirty two rats divided into four groups: 1) Control; 2) Compression ; 3) Compression + aquatic extract of *prosopis farcta* injections (50mgkg⁻¹ , ip ,2 time); 4) Compression + aquatic extract of *prosopis farcta* injections (75mgkg⁻¹ , ip ,2 time). (N=8)

Surgery

Animals were anesthetized under intera peritoneal injection of a mixture of ketamin and xylazine (6,66mgkg⁻¹).

Right sciatic nerve was exposed through a gluteal muscle splitting incision. At this location the nerve trunk was crushed for 60 seconds period between prongs of #5clamp forceps. The muscle and skin were then closed with 14mm stainless steel sutures (12). They could consume enough water and specified food during the experiment. In care groups, the extract injection was carried out immediately after compression during 28 days (Each week one injection).

After 28 day following perfusion a block of the spinal cord segments L4 to L6 (approximately 8mm length) was removed while sciatic nerve roots of both sides were still attached it. Since the nervous tissues are very sensitive and autolysis rapidly. Besides fixators cannot penetrate in spinal cord because of though cover around it. So for better fixation, perfusion method was used. When perfusion finished, sampling of spinal cord was began. The spinal cord was completely separated and 8mm samples at the injured area (L4-L6) entered to passage stage, then entered to cutting stage and serially 7 Mm sections were prepared and colored with toluidine blue. Required photos from front horn of spinal cord for future studies were taken according their numbers. Two photos were taken from two serial sections, one of anterior horn right half of first section and another from anterior horn right half of second section. The magnitude of microscope in this stage was $5\times 10\times 2/5=100$.

In order to count neuroglia cells random systematic method was used and dissector method was used for counting particles (13).

Statistical analysis

The ratio of numerical density of neuroglia cells in samples of spinal cord was used as an index of neuroglia cells death. All quantitative data were analyzed using ANOVA and t-test.

All data were presented as mean \pm S.E. Differences were considered statistically significant when ($p<0.05$).

Results

The results indicate several facts: 1-The number of neuroglia cells increases in compression group in compare with control group ($p < 0.05$).

2- The neuroglia cells density (number of neuroglia cells) in all experimental groups increased in compare with control group and very near to compression group (Table.1).

Table.1: Neuroglia cells density in anterior horn of spinal cord in different groups. Data are presented as mean \pm S.E. (n=8 in each group).

Groups	NV (Mean \pm S.E.)
Control	56 \pm 2
Compression	101 \pm 2
Treatment (50mgkg-1)	120 \pm 1
Treatment (75mgkg-1)	131 \pm 3

3- The neuroglia density in treatment groups is very similar with compression group.




Fig.1: Comparison of Neuroglia cells density in anterior horn of spinal cord in different groups. Data are presented as mean \pm S.E. (n=8 in each group).

A) The compression of the neuroglia cells density in treatment aquatic extract in 2 different dosages (50, 75 mg/kg) with compression group.

B) The compression of the neuroglia cells density in compression and control groups. ($p < 0.05$)*

4-The number and size of Neuroglia cells in experimental groups have remarkable increase (Fig.2).




Fig.2: The cross section of spinal cord (magnitude 200x And painting in toluidin blue)A: Alpha motoneuron B: Neuroglia cell

Discussion

Following injury Neuroglia cells were activated, microglial cells present a robust antigenic profile. They participate in the pathogenesis of neurological disorders by secreting various inflammatory molecules such as cytokines or nitric oxide. When CNS cells die, microglia can be further activated and become phagocytes. While considerably different in nature, all of these morphological and physiological properties have been used to identify microglial cells.

There are some evidences supporting the hypothesis that some herbs may also exert neurotrophic and inflammation actions (14). It provides neuronal differentiation and increase in neuritis outgrowth. This plants act by prevention of relieving glutamate (15).

Causing any pressure to axon or occurrence of axonic defeat induced changes in both distal and proximal parts because of breaking the connection of axon with neuron's body(16). In distal part, both axon and myelin cover is degenerated completely. In this defeat, emerging 2-3 days later, Endoneurim layers keep unchanged. Considering the ability of producing new myelin, Schwann cells increase along degenerated fiber, so in distal part of defeated nerve, for myelin making in new branches of proximal part of nerve new circumstances is required (17).

In these phenomena neuralgias proliferated and act as macrophage cells (18). In addition increasing of inflammation proceeding is coordinated with proliferated of neuroglia cells. Then if some component could inhibit these phenomena, degeneration of central nervous system was decreased.

So it is possible that *prosopis farcta* aquatic extract has protective effects on injured central nervous system cell and the results of this research show this hypothesis. It was shown that the neuroglia density in compression group increased in compare with control group (Table.1).The neuroglia density in treatments groups is very near to compression group. There is not any meaningful different between experimental and compression group in neuroglia density. It may be concluded pod aquatic extract of

prosopis farcta cannot protect neuroglia from proliferation after sciatic nerve injury. May be this component has had some roles in regeneration phenomena and act as a neurotrophic factors (Fig.2).

These factors consist of fibroblastic growth factors secreting from Schwann cells and macrophages that their synthesis is stimulated by cytokines and sticky molecules (19). Other neurotrophic factors affecting on correction are growth factor (NGF), growth factor originating from brain (BDNF), insulin – like growth factor (IGF) of integrin, laminin, collagen and fibronectin (20).

Following sciatic nerve injury, the generation of free radicals causes apoptosis in the cell body of spinal cord neurons (21). Antioxidants inactive free radicals and prevent apoptosis in spinal cord neurons. Different species of *prosopis farcta* have anti oxidant effects and are responsible for collection and removing free radicals (9). Therefore protective effects of aquatic extract are due to its anti- oxidant role.

In total administration of *prosopis farcta* pod aquatic extract after sciatic nerve injury cannot decrease neuroglia density in compare with compression groups.

Acknowledgment

This study was supported by Islamic Azad University of Mashhad, Iran. We thank Dr. Khayyatzade for her helpful comments and discussions.

References

1- Patro N, Nagayach A, Patro IK. Iba1 expressing microglia in the dorsal root ganglia become activated following peripheral nerve injury in rats. *Indian J Exp Biol*(2010);48(2):110-6.

2- Liu L, Rudin M, Kozlova EN. Glial cell proliferation in the spinal cord after dorsal rhizotomy or sciatic nerve transection in the adult rat. *Exp Brain Res* (2000);131(1):64-73.

3- Chaisuksunt V, Zhang Y, Anderson PN, Campbell G, Vaudano E, Schachner M, Lieberman AR. Axonal regeneration from CNS neurons in the cerebellum and brainstem of adult rats: correlation with the patterns of expression and distribution of messenger RNAs for L1, CHL1, c-jun and growth-associated protein-43. *Neuroscience*(2000);100(1):87-108.

4- Anderson PN, Campbell G, Zhang Y, Lieberman AR. Cellular and molecular correlates of the regeneration of adult mammalian CNS axons into peripheral nerve grafts. *Prog Brain Res*(1998);117:211-32

5- Gordon T. The physiology of neural injury and regeneration: The role of neurotrophic factors. *J Commun Disord*(2010);43(4):265-73.

6- Vanelderden P, Rouwette T, Kozicz T, Roubos E, Van Zundert J, Heylen R, Vissers K. The role of brain-derived neurotrophic factor in different animal models of neuropathic pain. *Eur J Pain*. (2010);14(5):473.e1-9.

- 7- Cusick CG. Extensive cortical reorganization following sciatic nerve injury in adult rats versus restricted reorganization after neonatal injury: implications for spatial and temporal limits on somatosensory plasticity. *Prog Brain Res* (1996);108:379-90.
- 8- Deputy."The Iranian Plant's Pharmacopoeia." Iranian Ministry of Health and Medical Education, Deputy of Food and Medication (report) (2002) pp,1-3.
- 9- Al-Mustafa AH, Al-Thunibat OY. Antioxidant activity of some Jordanian medicinal plants used traditionally for treatment of diabetes. *Pak J Biol Sci*(2008) ;11(3):351-8.
- 10- Soares De Moura R, Costa Viana FS, Souza MA, Kovary K, Guedes DC, Oliveira EP, Rubenich LM, Carvalho LC, Oliveira RM, Tano T, Gusmão Correia ML. Antihypertensive, vasodilator and antioxidant effects of a vinifera grape skin extract. *J Pharm Pharmacol*(2002);54(11):1515-20.
- 11- Cicchetti E, Chaintreau A. Comparison of extraction techniques and modeling of accelerated solvent extraction for the authentication of natural vanilla flavors. *J Sep Sci* (2009);32(11):1957-64.
- 12- Behnam-rasouli M, Nikravesh M, Mahdavi M, Tehranipour M. Post-operative time effects after sciatic nerve crush on the number of alpha motoneurons, using a stereological counting method(Disector). *Iranian biomedical journal*(2000);4(1):45-49.
- 13- Tehranipour M, Kabiri M. The effect of exogenous testosterone administration on peripheral nerves regeneration after sciatic nerve compression in Rat. *J.B.S.*(2009)9(7):692-696.
- 14- Perry N, Court G, Bidet N, Court J, Perry E .European herbs with cholinergic activities: potential in dementia therapy. *Int J Geriatr Psychiatry* (1996);11:1063-1069.
- 15- Maier IC, Schwab ME. Sprouting, regeneration and circuit formation in the injured spinal cord: factors and activity. *Philos Trans R Soc Lond B Biol Sci*(2006);29:361(1473):1611-34.
- 16- Sakurai M, Nagata T, Abe K, Horinouchi T, Itoyama Y, Tabayashi K. Survival and death-promoting events after transient spinal cord ischemia in rabbits: induction of Akt and caspase3 in motor neurons. *J Thorac Cardiovasc Surg* (2003);125(2):370-7.
- 17- Rosenfield J, Paksima N. Peripheral nerve injuries and repair in the upper extremity. *Bull Hosp Jt Dis* (2001);60(3-4):155-61.
- 18- Hu P, McLachlan EM. Macrophage and lymphocyte invasion of dorsal root ganglia after peripheral nerve lesions in the rat. *Neuroscience*(2002);112(1):23-38.
- 19- Zhang Y, Roslan R, Lang D, Schachner M, Lieberman AR, Anderson PN. Expression of CHL1 and L1 by neurons and glia following sciatic nerve and dorsal root injury. *Mol Cell Neurosci*(2000);16(1):71-86.

20- Imaizumi, T., Lankford, K.L. and Kocsis, J.D. Transplantation of olfactory ensheathing cells or Schwann cells restores rapid and secure conduction across the transected spinal cord. *Brain Res*(2000);854(1-2),70-8.

21-Siniscalco D, Fuccio C, Giordano C, Ferraraccio F, Palazzo E, Luongo L, Rossi F, Roth KA, Maione S, de Novellis V. Role of reactive oxygen species and spinal cord apoptotic genes in the development of neuropathic pain. *Pharmacol Res* (2007);55(2):158-66