

ANTI-INFLAMMATORY AND ANALGESIC ACTIVITIES OF ETHANOLIC EXTRACT OF GLOCHIDION THOMSONII (BARK)

Md. Habibur Rahman¹, Abul Bassar Md. Oliullah¹, Md. Mizanur Rahman¹; Md. Jahir Alam², Masum Shahriar², Farhana Israt Jahan¹

¹Department of Pharmacy, Daffodil International University, Dhaka, Bangladesh.

²Department of Pharmacy, Jahangirnagar university, Saver. Dhaka, Bangladesh

Email address: mizanur.ph@diu.edu.bd, jifarhana@daffodilvarsity.edu.bd

Abstract

Objective: The present communication attempts to evaluate the Anti-inflammatory and Analgesic activities of ethanolic extract of *Glochidion Thomsonii* (Bark) belongs to the family: Euphorbiaceae, which is widely distributed in Bangladesh and India.

Methods: Ethanolic extract of *Glochidion Thomsonii* at doses of 250mg/kg and 500mg/kg were tested for their anti-inflammatory activity against Carrageenan induced paw edema & Xylene-Induced Ear Edema test in rats and analgesic activity was tested against Formalin-induced Paw licking test & acetic acid induced writhing (in mice) models and the two phases of formalin induced paw licking models of pain.

Results: The results indicate that the ethanolic extracts at the doses of 250 and 500mg/kg body weight significantly inhibited, the carrageenan induced paw edema and Xylene-induced ear edema models of inflammation, similarly the extract at the doses of 250 and 500mg/kg significantly inhibited the acetic acid induced writhing (in mice) models and the two phases of formalin induced paw licking models of pain.

Conclusion: This study confirms that the ethanolic extract of *Glochidion Thomsonii* has good analgesic and anti-inflammatory property. Further studies are necessary to isolate the compounds from the ethanolic extract of *Glochidion Thomsonii* responsible for the activity.

Keywords: *Glochidion Thomsonii*, analgesic, inflammation, extracts, rats and mice

Introduction

From the ancient time plants have been used in medicinal preparations for treatment of various human and animals diseases. Reduced efficacy of synthetic preparations due to various reasons leads the global interest in the preparation of therapeutic medicine from plants [1]. According to the world health organization (WHO), 70-80% of world population uses the plant derived drugs for treatment of various health problems [2].

Glochidion thomsonii (bark) belonging to the Euphorbiaceae in English *Phyllanthus thomsonii*, local name: Latikath, Bread flower, is a tall climbing shrub which is locally used as medicinal plant. It is distributed in Tropical Asia, Pacific islands and Malaysia. A few in Tropical America and Africa and also found in Bangladesh. The plants having triterpenoids are the most widely used for the treatment of inflammation and many other life threatening diseases in the traditional medicine of different cultures [3], In India a number of medicinal plants and their formulations are widely used for the treatment of various disorders[4]. Therefore in the present study we aim to investigate whether the ethanolic extract of *Glochidion thomsonii* (bark) has analgesic and anti-inflammatory activities.

Methods

Animals

For the experiment Swiss albino mice of either sex, 6-7 weeks of age, weighing between 20-30gm were collected from the animal research lab in the Department of Pharmacy Jahangirnagar University, Savar, Dhaka. Animals were maintained under standard environmental conditions (temperature: $27.0 \pm 1.0^\circ$), relative humidity: 55-65% and 12 h light/12 h dark cycle) and had free access to feed and water ad libitum. The animals were acclimatized to laboratory condition for one week prior to experiments. All protocols for animal experiment were approved by the institutional animal ethical committee.

Plant Materials

The bark of plant of *Glochidion thomsonii* was collected from Sylhet forest area during January 2017. The plant material was taxonomically identified by the National herbarium of Bangladesh. The plant

material was taxonomically identified by the National herbarium of Bangladesh.

Preparation of Plant Extract

The plant material was shade-dried with occasional shifting and then powdered with a mechanical grinder, passing through sieve #40, and stored in a tight container. About 500 gm of powdered sample was taken in a clean, flat-bottomed glass container and soaked in 4000 ml of 96% ethanol. The container with its contents was sealed and kept for a period of 10 days accompanying occasional shaking and stirring. The whole mixture then underwent a coarse filtration by a piece of clean, white cotton material. Then it was filtered through whatman filter paper. The filtrate was kept in an open space to evaporate the solvent thus crude extract was obtained.

The collected mixture of active constituents with ethanol was dried with a Rota evaporator (BUCHI Rota Vapor R-114, Switzerland) under reduced pressure to get viscous substance. Then it was transferred to a beaker and taken on a water bath for further drying at room temperature. Finally a solid mass was obtained and preserved in a Petridis in the refrigerator. Extract obtained was 49 gm from 500 gm powder using 4000 ml ethanol. Yield = 9.80% w/w.

Phytochemical analysis

Desirable amount of *Glochidion thomsonii* bark extract was solubilized in water for phytochemical analysis. The extracted solution was tested for alkaloids, glycosides, steroids, gums, flavonoids, saponins, sugars and tannins according to the protocol described by Trease and Evans [5].

Analgesic activity

Formalin-induced Paw licking test

The method of Hunskaar and Hole (1987) was used for the study. Mice were divided into 4 groups of 6 animals each. Group I, the control group received distilled water, group II, the standard group received Indometacin BP (100 mg/kg). Groups 3 and 4 received alcoholic extract (250 mg/kg and 500 mg/kg). After 1 hour of drug administration, 2.7% formalin was injected into the dorsal surface of the left hind paw. The time spent licking the injected paw was recorded. Animals were observed for the 5 min post

formalin (acute phase) and for 5 min starting at 20th min post formalin (delayed phase) (Hunskaar and Hole, 1987) [6].

Acetic acid induced writhing test

For analgesic test all mice were divided into four groups. Each group comprises of 5 mice. First group are controlled group here used to distilled water. Group two are standard group here used to Indometacin BP (100mg/kg). Group three and four received to alcoholic extract (250 & 500 mg/kg). 45 minutes later each mouse was injected with 0.7% acetic acid at a dose of 10 mL/kg body weight. The number of writhing responses was recorded for each animal during a subsequent 3 min period after 15 min of the I.P. administration of Acetic acid and the mean abdominal writhes for each group was obtained (Koster, et al., 1959).[7].

The percentage inhibition was calculated using the formula:

$$\text{Inhibition(\%)} = \frac{\left(\frac{\text{Mean number of writhes (control)}}{\text{Mean number of writhes (control)}} \right) - \left(\frac{\text{Mean number of writhes (drugs)}}{\text{Mean number of writhes (control)}} \right)}{\left(\frac{\text{Mean number of writhes (control)}}{\text{Mean number of writhes (control)}} \right)}$$

Anti-inflammatory action evaluation

Carrageenan induced paw edema

Pedal inflammation was produced in rats according to the carrageenan induced paw edema method [8]. 24 albino rats were divided into four groups of six each and fasted overnight for 18 hrs with water. Next day the animals were weighted and numbered. A mark was made on the right hind paw just beyond tibio-tarsal junction, so that every time the paw was dipped in the mercury (Hg) column up to the fixed mark to ensure constant paw volume. The initial paw volume was noted of each rat by mercury (Hg) displacement method. 0.1 ml of 1 % carrageenan was injected into the right hind paw of each rat under the sub plantar aponeurosis. The animals were treated orally with extracts (250 and 500 mg/kg), indomethacin (100mg/kg) or Saline (10 ml/kg) 1 h before carrageenan injection. The paw volume of each rat after carrageenin injection after one hour, two hour, three hour and five hour was recorded by mercury displacement in plethysmograph.

This volume is called final volume. The anti-inflammatory activity of the extract was measured by

its potential to prevent edema caused by carrageenan as against the control group which was given the vehicle only. Mean paw edema was calculated for each group as average of paw volume of individual rats belonging to that group. Since, the mean was subjected to positive and negative fluctuations hence; standard error for each group was also calculated.

$$\text{Standard error (S.E)} = \text{standard deviation} / \sqrt{n}$$

Where, n = number of rats in each group

Percent inhibition of paw edema was calculated according to the following formula:

$$\% \text{ Inhibition} = a-b/a \times 100$$

a- is the mean paw inflammation volume of control.

b- is the mean paw inflammation volume of test

Xylene-Induced Ear Edema test

The Xylene-induced ear edema test was performed as described in Dai et al. (1995). Mice were divided into 4 groups of 6 animals each. Group I, the control group received distilled water. Group II, the standard group received Indomethacin [100 mg/kg]. Groups III and IV received alcoholic extract (250 mg/kg and 500 mg/kg). One hour later, each animal received 20µl of xylene on the anterior and posterior surfaces of the right ear lobe. The left ear was considered as control. Mice were sacrificed one hour after Xylene application and circular sections were taken, using a cork borer with a diameter of 3 mm, and weighed. The percentage of ear edema was calculated as inflammation based on the weight of left ear without xylene (Dai et al., 1995)[9].

Results

The results manifest the presence of glycoside, steroids, flavonoids, tannin, saponins and reducing sugars from ethanolic crude extract of the *Glochidion thomsonii* bark (see Table 1).

Analgesic Studies

Formalin-induced paw licking

The ethanolic extract of *Glochidion thomsonii* at the doses of 250 and 500 mg/kg significantly inhibited the two phases of the formalin test. The summary of the result are shown in Table 2.

Acetic acid induced writhing test

The ethanolic extract of *Glochidion thomsonii* at the doses of 250 and 500 mg/kg significantly decreased the number of writhing from 11.40 ± 2.94 to 10.50 ± 2.90 . The summary of the results are shown in Table 3.

Anti-Inflammatory Studies

Carrageenan induced paw edema

The ethanolic extract of *Glochidion thomsonii* at the doses of 250 and 500 mg/kg significantly reduced the carrageenan induced edema. The results are shown in Table 4.

Xylene-Induced Ear Edema test

The ethanolic extract of *Glochidion thomsonii* at the doses of 250 and 500 mg/kg significantly inhibited the two phases of xylene induced inflammation. The summary of the result are shown in Table 5.

Discussion

The results obtained from the present study shows that the ethanolic extract of *Glochidion thomsonii* has analgesic and anti-inflammatory activities. The extracts significantly inhibited the licking time in the two phases of the formalin test as well as the characteristic writhing observed following intraperitoneal injection of acetic acid. These two tests (formalin induced paw licking and acetic acid induced writhing) are used for detecting specific activities of drugs that may have analgesic activities. Drugs that inhibit the first phase of the formalin test have the ability to alleviate neurogenic pain while those drugs that inhibit the second phase of the test have the ability to inhibit inflammatory pain [10]. The acetic acid induced writhing test on the other hand is a model of visceral pain that is useful for analgesic drug development except that it gives false positives for muscle relaxant and sedatives [11,12] Since the extract of *Glochidion thomsonii* produced significant effects in all the models of pain used in this study it show that the extract at high doses has strong analgesic activity. The anti-inflammatory activity of the ethanolic extract of *Glochidion thomsonii* was investigated using two models namely; carrageenan, and Xylene-induced ear edema test models which represent acute and chronic forms of inflammation

[13]. The extract dose dependently inhibited both two types of inflammation although it was the higher doses of the extracts that were highly effective against both the two types of inflammation Above extract at concentration of 500mg/kg was found to inhibit 94.4% of edema measured after 4 hrs of injecting 0.1ml of 1% (w/v) of carrageenan, which was comparable to Indomethacin [100 mg/kg] taken as standard, showing 88.9% of edema inhibition after 4 hrs. A marked decrease in mean paw edema was observed in both the extract treated groups observed at 1hr, 2hr, and 3hr and 4hr favoring it was more explicit in the higher dose (500mg/kg) group.

Conclusion

In conclusion, this study demonstrated that the ethanolic extract of *Glochidion thomsonii* contain Glycoside, alkaloid, flavonoid, saponin, tannin and steroid type of compounds and have a significant analgesic and anti-inflammatory activity. Further studies will be necessary to identify the possible compounds responsible for the anti-inflammatory and analgesic activities of different extracts of *Glochidion Thomsonii*.

Acknowledgments

Authors are thankful to Daffodil International University and Jahangirnagar University for providing the necessary lab space and equipment for this research work.

References

1. Burkill HM. The useful plants of west tropical Africa. Volume 2: Families El. Royal Botanic Gardens; 1994.
2. Bastos GN, Santos AR, Ferreira VM, Costa AM, Bispo CI, Silveira AJ, Do Nascimento JL. Antinociceptive effect of the aqueous extract obtained from roots of *Physalis angulata* L. on mice. *Journal of ethnopharmacology*. 2006 Jan 16;103(2):241-5.
3. Divakar CM, Sheela S, Sandhya S, Vinod KR, Pillai NR, Rao SB. Anti-inflammatory and Antioxidant activities of *Polyscias filicifolia* saponins. *Der Pharmacia Lettre*. 2010;2(1):41-7.
4. Bhaskar VH, Balakrishnan N. Hepatoprotective activity of laticiferous plant species (*Pergularia daemia* and *Carissa carandas*) from Western Ghats,

Tamilnadu, India. Der Pharmacia Lettre. 2009;1(2):130-42.

5. Trease GE and Evans WC: Pharmacognosy, 1st Edn. London: Brailliere Tindall Press. 1983; 307-706

6. Hunskaar S, Hole K. The formalin test in mice: dissociation between inflammatory and non-inflammatory pain. Pain. 1987 Jul 1;30(1):103-14.

7. Koster R. Acetic acid for analgesic screening. InFed proc 1959 (Vol. 18, p. 412).

8. Winter CA, Risley EA, Nuss GW. Carrageenin-induced edema in hind paw of the rat as an assay for antiinflammatory drugs. Proceedings of the society for experimental biology and medicine. 1962 Dec;111(3):544-7.

9. Dai Y., Liu L H., Kou J P. J. China Pharm. Uni., 1995; 6: 362—364.

10. Tjølsen A, Berge OG, Hunskaar S, Rosland JH, Hole K. The formalin test: an evaluation of the method. Pain. 1992 Oct 1;51(1):5-17.

11. Elisabetsky E, Amador TA, Albuquerque RR, Nunes DS, do CT Carvalho A. Analgesic activity of *Psychotria colorata* (Willd. ex R. & S.) Muell. Arg. alkaloids. Journal of Ethnopharmacology. 1995 Oct 1;48(2):77-83.

12. Vyklícky L. Techniques for the study of pain in animals. Advances in pain research and therapy. 1979;3:727-45.

13. Ismail TS, Gopalakrishnan S, Begum VH, Elango V. Anti-inflammatory activity of *Salacia oblonga* Wall. and *Azima tetraantha* Lam. Journal of ethnopharmacology. 1997 Apr 1;56(2):145-52.

Table 1: phytochemical properties of *Glochidion thomsonii* crude extract

Compounds	Observation
Glycosides	+ ve
Alkaloids	- ve
Steroids	+ ve
Gums	- ve
Flavonoids	+ ve
Saponins	+ ve
Reducing sugars	+ ve
Tannins	+ ve

Table 2: Effects of the ethanolic extract of *Glochidion thomsonii* on Formalin-Induced Paw Licking Test

Groups	No. of animal	Dose (mg/kg b.wt.)	Licking time(sec)a	
			Early phase	Late phase
Control	06	10 ml/kg	131.6 ± 5.8	89.60 ± 5.8
Indometacin BP	06	100 mg/kg	68.20 ± 5.4**	27.20 ± 5.5**
Extract I	06	250 mg/kg	64.00 ± 3.7	35.60 ± 7.9
Extract II	06	500 mg/kg	52.80 ± 4.6***	26.20 ± 2.4***

Each value is the mean ± S.E.M. of 6 rats

*P < 0.05; **P < 0.01; ***P < 0.01 compared with control; student's t-test.

Table 3: Effect of the ethanolic extract of *Glochidion thomsonii* on Acetic acid induced writhing in mice

Groups	No. of animals	Dose (mg/kg b.wt)	Writhing count	% of inhibition of analgesia
Control	06	10 ml/kg	23.80 ± 0.92	00
Indomethacin	06	100 mg/kg	8.40 ± 1.08**	64.71
Extract I	06	250 mg/kg	11.40 ± 2.94*	52.10
Extract II	06	500 mg/kg	10.50 ± 2.90*	55.88

Each value is the mean ± S.E.M. of 6 rats

*P < 0.05; **P < 0.01; ***P < 0.01 compared with control; student's t-test

Table 4: Effects of ethanolic extract of *Glochidion thomsonii* on Carrageenan-induced paw

Group	No. of animal	Dose (mg/kg.wt.)	Mean increase in paw volume		Inhibition (%)	
			3hr	4hr	3hr	4hr
Control	06	10 ml/kg	3.6 ± 0.60	3.4 ± 0.75		
Indomethacin	06	100 mg/kg	1.6 ± 0.24*	0.4 ± 0.45*	55.6	88.9
Extract I	06	250 mg/kg	0.8 ± 0.20**	0.8 ± 0.20*	77.8	77.8
Extract II	06	500 mg/kg	0.6 ± 0.24**	0.2 ± 0.20*	83.3	94.4

*Each value is the mean ± S.E.M. of 6 rats

*P < 0.05; **P < 0.01; ***P < 0.01 compared with control; student's t-test.