

ASCORBIC ACID CONTENT IN THE HERBAL MIXTURE WITH ANTIDIABETIC ACTIVITY

Savych, Alona^{*1}, Basaraba, Roksolana²

¹Department of Pharmacognosy with Medical Botany, I. Horbachevsky Ternopil National Medical University, Ukraine

²Department of Pharmacy, Bukovina State Medical University, Ukraine

*alonasavych@gmail.com

Abstract

Diabetes mellitus is a global social problem in the field of health care, due to rapid spread of this disease and the development of serious complications such as micro- and macroangiopathies, which significantly reduce the quality and life expectancy of patients.

The aim of the research was to study ascorbic acid content in the herbal mixture (Equiseti arvensis herba, Sambuci flores, Inulae rhizomata et radices, Hyperici herba, Tiliae flores, Polygoni avicularis herba, Myrtilli folium, Urticae folia), which have established hypoglycemic, hypolipidemic and antioxidant activity in previous studies in vivo and to validate the method that was used.

The study of vitamin C content in herbal sample was carried out by spectrophotometry (λ_{max} 520 nm) with the measurement of the absorption spectrum in terms of L-ascorbic acid. The method used was validated for linearity, sensitivity, precision and was established limit of detection and limit of quantification.

According to the results of spectrophotometric determination, it was found that ascorbic acid content in the herbal mixture was 0.13 ± 0.02 %.

The spectrophotometric method for the analysis of ascorbic acid content was validated and showed good linearity of the standard solution of L-ascorbic acid (0.02-0.1 mg/mL). The calibration curve was expressed by the regression equation $y = 0.0371x - 0.0081$, the correlation coefficient was $R^2 = 0.9929$, the limit of detection – 0.01 mg/mL, and limit of quantification – 0.04 mg/mL. The method used meets the characteristics of precision and accuracy.

The data obtained indicate the presence of a correlation between the phytochemical composition of the studied herbal mixture and its pharmacodynamics, which was previously established.

Keywords: *herbal mixture, ascorbic acid, vitamin C, UV-spectrophotometry, diabetes mellitus, phytotherapy, antioxidant activity, validation*

Introduction

Diabetes mellitus is a global social problem in the field of health care, due to rapid spread of this disease and the development of serious complications such as micro- and macroangiopathies, which significantly reduce the quality and life expectancy of patients [1, 2]. According to the official information of International Diabetes Federation (2019) the number of patients is projected to increase to 642 million by 2040 [3]. An important problem of pharmacovigilance is that existing pharmacotherapy is able to reduce hyperglycemia, but it is not always able to stabilize fluctuations of glycemia values during the day and keep it at optimal level [4, 5, 6]. This leads to the formation of a cascade of pathological processes – excessive glycation and inactivation of the body's antioxidant defense system, triggering free radical oxidation of lipids and, as a consequence, the development of oxidative stress, which leads to the development and progression of diabetic complications [7, 8]. In turn, oxidative stress, which is the result of the accumulation of reactive oxygen species (ROS), reactive nitrogen molecules (RNS) and some derivatives of heavy metals are the main event for the development of insulin resistance [8, 9]. This may reduce peripheral insulin sensitivity through major molecular mechanisms such as β -cell dysfunction, inflammatory responses, decreased regulation and / or localization of glucose transporter 4 (GLUT-4), mitochondrial dysfunction, and abnormal insulin signaling pathways [1, 4, 11]. Therefore, the optimization of pharmacotherapy, search and study of new remedies with antioxidant activity for the prevention and treatment of this disease and its dangerous complications is a top issue of pharmacy and medicine.

One of these areas is phytotherapy, as it has a number of advantages over traditional therapy with using oral synthetic agents, namely, it is low-toxic, has a mild pharmacological effect and possibility to be used for long periods of time without significant side effects, is well combined with synthetic drugs, has a complex activity through a number of biologically active compounds [12, 13, 14, 15]. Particular attention deserve the combinations of different medicinal plants [16, 17], because such plant mixtures are expected to have more

biologically active substances that influence on all links of the pathogenetic mechanism of development of diabetes mellitus and its complications [18, 19]. Plant biocompounds have a wide range of pharmacological action and a variety of mechanisms of influencing on the development of diabetes and diabetic angiopathies [4, 9].

In this regard, the important biologically active substance is ascorbic acid, namely vitamin C. Therefore, the aim of study was to investigate an ascorbic acid content in herbal mixture (*Equiseti arvensis herba, Sambuci flores, Inulae rhizomata et radices, Hyperici herba, Tiliae flores, Polygoni avicularis herba, Myrtilli folium, Urticae folia*) with previously studied antidiabetic activity *in vivo* [20, 21, 22] and validate this method of determination.

Methods

Plant materials: It was used the herbal raw materials harvested in June – August 2017 in Ternopil region and Carpathians (*Vaccinium myrtillus* leaf) (Ukraine) during the study. After harvesting, the raw materials were dried, crushed and brought back to standard according to the general GACP requirements [23]. The plants were identified by Department of Pharmacognosy with Medical Botany, I.Horbachevsky Ternopil National Medical University, Ternopil, Ukraine. Samples of herbal raw materials have been deposited in Departmental Herbarium for future record.

Chemicals and standards: United States Pharmacopeia (USP) reference standard of L-ascorbic acid was of primary reference standard grade ($\geq 99\%$ purity HPLC) and was purchased from Sigma-Aldrich Chemical Company (Germany). Oxalic acid, methanol, ethanol, thiourea, 2,6-dichloroindophenol sodium salt, 2,4-dinitrophenylhydrazine (2,4-DNPH), sulfuric acid was purchased from Sfera Sim LTD (Ukraine). Water used in the studies was produced by MilliQ Gradient water deionization system (USA).

Instrumentation and conditions: spectrophotometric measurements were performed on spectrophotometer UV-1800 Shimadzu (Japan) equipped with 1 cm quartz cells. Determination of ascorbic acid content was performed at a maximum absorption of 520 nm [23].

Reagents: Reagent A – 1.00 g of oxalic acid was

dissolved in 50 mL of methanol. *Reagent B* – 100.0 g of thiourea was dissolved in 1 L of ethanol 50% (v/v). *Reagent C* – 1.5 g of 2,4-DNPH was dissolved in 50 mL of 20 % solution of sulfuric acid. *Indicator* – 50.0 mg of 2,6-dichloroindophenol sodium salt was dissolved in 100 mL of distilled water.

Test solution: 1.00 g (accurately mass) of each powdered herbal raw materials was placed into a round-bottomed flask. Then was added *reagent A* and heated on a water bath under reflux for 10 minutes. After that, it was cooled in an ice bath to a temperature of (15–20) ° C and filtered through a paper filter. Then 2 mL of the filtrate was transferred into a 50 mL conical flask, added sequentially, gently shaking after each addition of 2 mL of an *indicator*. Subsequently, exactly 60 s later, 0.5 mL of *reagent B* and 0.7 mL of *reagent C* were added, heated under reflux at 75 ° C and immediately placed in an ice bath for 5 minutes. Then 5 mL of a mixture consisting of 12 mL of water and 50 mL of concentrated sulfuric acid was added dropwise to the obtained solutions, adding for a period of not less than 90 s and not more than 120 s, vigorously shaking the flask in an ice bath. Maintained for 30 min at room temperature and measured the optical density using *reference solution A* as a compensating solution.

Reference solution A: 2 mL of the filtrate obtained in the preparation of the test solution was treated as described above by adding *reagent C* immediately before measuring the optics.

Standard solution: 40.0 mg (accurately mass) of USP of ascorbic acid was dissolved in 100 mL of *reagent A*. Then 5 mL of obtained solution was placed into 100 mL volumetric flask and made up to the mark by *reagent A*. 2 mL of the resulting solution was worked up as described above for the filtrate obtained in the preparation of the test solution. The optical density was measured using *reference solution B* as a compensating solution.

Reference solution B: 2 mL of the standard solution was treated as described above by adding *reagent C* immediately before measuring the optics.

Method validation: the method was validated for linearity, sensitivity and precision (intra-assay and intermediate precision), accuracy, LOD, and LOQ [24].

Calibration curve and linearity: standard solution of ascorbic acid (2 mg/mL) was prepared in two replicate and diluted to concentrations of 0.02 mg/mL, 0.04 mg/mL, 0.06 mg/mL, 0.08 mg/mL and 0.1 mg/L. The calibration curve was constructed by plotting the measurements of mean absorbance versus the concentration of the standard solutions. The results were analyzed by linear regression and the correlation coefficient (R^2).

Sensitivity: the values for limit of detection (LOD) and limit of quantitation (LOQ) were calculated based on the data obtained during linearity testing in the low concentration range of the working in the test solution, using the following formulas: $LOD = 3.3 * s / \text{Slope}$; $LOQ = 10 * s / \text{Slope}$.

Precision and accuracy: precision was determined through the determination of intraassay and intermediate precision (41). Intra-assay precision was determined by measuring three different concentrations of each standard (prepared as two replicates, each measured three times) in one laboratory by one person. Intermediate precision was carried out by measuring the same concentrations of standards (prepared as two replicates, each measured three times) by two people in the same laboratory. Precision was expressed as percentage coefficient of variation (CV), using the following formulas: $CV = \text{standard deviation} / \text{mean} \times 100$. Accuracy is expressed through recovery, which was determined by measuring known concentrations of standard (in three concentration levels, in triplicate).

Statistical Analysis MS Excel (Microsoft, Redmond, WA) was used for data analysis. The results for the calibration curves were based on two replicate samples of each standard concentration, each measured twice ($n = 4$). Regression was used to obtain the linear equation and determine R^2 . Precision and recovery studies were based on the measurement of three different concentration levels, each as two replicates and each measured three times ($n = 6$). Ascorbic acid content in the herbal mixtures was the results of two replicate samples, each measured twice ($n = 4$) and were presented as mean value \pm SD.

Results and Discussion

The results of spectrophotometric determination

of the ascorbic acid content are presented in Figure 1. The method of determination was based on conversion of an ascorbic acid to coloured complex with 2,4-DNPH. The maximum absorption of all samples was at $\lambda_{\max} = 520 \pm 0.5$ nm.

According to spectrophotometric determination, it was found that ascorbic acid content was 0.13 ± 0.02 % in the herbal mixture (Fig. 1).

The analytical procedure has been validated to confirm its reliability. Calibration solutions were prepared in triplicate and analyzed under the optimal conditions as described above. The calibration curve was found to be linear in the range 0.02-0.1 mg/mL for USP of ascorbic acid (Fig. 2). Regression equation ($y=0.0371x-0.0081$) and $R^2=0.9929$ revealed a good linearity response for spectrophotometric method developed for the determination of an ascorbic acid content.

The results of validation showed that the LOD and the LOQ of USP of ascorbic acid was 0.01 mg/mL and 0.04 mg/mL, respectively, indicating that the sensitivity of the method was satisfactory (Table 1).

In the table 2 represents the precision and accuracy of spectrophotometric method for the determination of ascorbic acid content. CVs for the spectrophotometric method were between 1.7 % and 3.2 % and between 1.8 and 4.5 % for intra-assay and intermediate precision, respectively. Precision was studied through intra-assay and intermediate precision. Accuracy was expressed through the determination of recovery, which was between 97.2 % and 105.1 %.

The incidence of diabetes type 2 is increasing worldwide, and patients with diabetes have a greatly increased risk of death from cardiovascular disease [25]. This risk may be associated with hyperglycemia, insulin resistance, dyslipidemia and hypertension. In addition, an important pathogenetic significance in the development of cardiovascular complications in patients with diabetes mellitus type 2 is an increase in arterial stiffness. Literary data indicate that ascorbic acid has the ability to improve vascular elasticity, which in turn will restore microcirculation [26].

Ascorbic acid, an antioxidant vitamin, plays an important role in protecting free radical-induced damage. Antioxidant activity of vitamin C in the treatment and prevention of diabetes and its

complications is important because they can include suppression of ROS formation either by inhibition of enzymes or by chelating trace elements involved in free radical generation [27]; scavenging ROS; inhibition the enzymes involved in ROS generation – microsomal monooxygenase, glutathione S-transferase, mitochondrial succinoxidase, nicotinamide adenine dinucleotide phosphate (NADH) oxidase, and so forth [28].

Literature sources indicate that diabetes mellitus type 2 reduces the level of basal vitamin C. Vitamin C is structurally similar to glucose and can replace it in many chemical reactions and thus is effective for the prevention of nonenzymatic glycosylation of protein [27, 28].

Thus, the high content of ascorbic acid in the studied herbal mixture (Fig. 1) may indicate the ability of this mixture to prevent the development of oxidative stress, which is the main pathogenic mechanism of diabetic angiopathies, and can prevent the development of cardiovascular disease [29]. In addition, the obtained results are reliable because the method to determine ascorbic acid content in the studied object was validated for linearity, sensitivity and precision (intra-assay and intermediate precision), accuracy, LOD, and LOQ (Fig. 2, Table 1-2).

The obtained data testify to the expediency of using the studied herbal mixture in order to optimize antidiabetic pharmacotherapy.

Conclusions

The results of spectrophotometric analysis indicate the high content of ascorbic acid in the studied herbal mixture that provides the powerful antioxidant activity, which is an important factor in the treatment of diabetes and in preventing the development of diabetic angiopathies. The ascorbic acid content was 0.13 ± 0.02 % in the herbal mixture, which contains *Equiseti arvensis herba*, *Sambuci flores*, *Inulae rhizomata et radices*, *Hyperici herba*, *Tiliae flores*, *Polygoni avicularis herba*, *Myrtilli folium*, *Urticae folia*.

Spectrophotometric method to determine vitamin C content was validated and showed good linearity of the standard solution of L-ascorbic acid in the range of 0.02-0.1 mg/mL, high sensitivity of the method, precision (intra-assay and intermediate

precision) and accuracy.

The obtained phytochemical studies may indicate a correlation between the component composition and content of ascorbic acid in the sample of studied mixture and its effectiveness in the treatment and prevention of diabetes mellitus and its complications.

References

- American Diabetes Association (2020). Standards of Medical Care in Diabetes. *Diabetes care*, 43, 1212.
- Savych, A. O., & Marchyshyn, S. M. Phytochemical composition of the herbal mixtures, which causes their effectiveness in diabetes mellitus. *Challenges and achievements of medical science and education: Collective monograph*. Riga, Latvia: "Baltija Publishing", 276-290.
- International Diabetes Federation (2019). *IDF Diabetes Atlas, 9th ed.* Brussels, Available at: <https://www.diabetesatlas.org>
- Skyler, J. S., Bakris, G. L., Bonifacio, E., Darsow, T., Eckel, R. H., Groop, L., Groop, P. H., Handelsman, Y., Insel, R. A., Mathieu, C., McElvaine, A. T., Palmer, J. P., Pugliese, A., Schatz, D. A., Sosenko, J. M., Wilding, J. P., & Ratner, R. E. (2017). Differentiation of Diabetes by Pathophysiology, Natural History, and Prognosis. *Diabetes*, 66(2), 241–255.
- Savych, A., Marchyshyn, M., Basaraba, R., & Lukanyuk, M. (2020). Antihyperglycemic, hypolipidemic and antioxidant properties of the herbal mixtures in dexamethasone-induced insulin resistant rats. *PharmacologyOnLine*, 2, 73-82.
- Savych, A., Marchyshyn, M., & Naconechna, S. (2021). Influence of some herbal mixtures on insulin resistance and glucose tolerance in rats. *PharmacologyOnLine*, 1, 356-364.
- Ndjaboue, R., Farhat, I., Ferlatte, C. A., Ngueta, G., Guay, D., Delorme, S., Ivers, N., Shah, B. R., Straus, S., Yu, C., & Witteman, H. O. (2020). Predictive models of diabetes complications: protocol for a scoping review. *Systematic reviews*, 9(1), 137.
- Savych, A., Marchyshyn, M., & Basaraba, R. (2020). Screening study of hypoglycemic activity of the herbal mixtures (Message 1). *ScienceRise: Pharmaceutical Science*, 4(26), 40-46.
- Oguntibeju O. O. (2019). Type 2 diabetes mellitus, oxidative stress and inflammation: examining the links. *International journal of physiology, pathophysiology and pharmacology*, 11(3), 45–63.
- Choi, S. W., & Ho, C. K. (2018). Antioxidant properties of drugs used in Type 2 diabetes management: could they contribute to, confound or conceal effects of antioxidant therapy? *Redox report: communications in free radical research*, 23(1), 1–24.
- Savych, A., Marchyshyn, S., Kyryliv, M., & Bekus, I. (2021). Cinnamic acid and its derivatives in the herbal mixtures and their antidiabetic activity. *Farmacia*, 69(3), 595-601.
- Marchyshyn, S., Polonets, O., Savych, A., & Nakonechna, S. (2020). Determination of carbohydrates of *Chrysanthemum morifolium* L. leaves and flowers by GC-MS. *Pharmakeftiki*, 32(4), 202-212.
- Savych, A.O., Marchyshyn, S.M., Kozyr, H.R., & Skrinchuk, O.Y. (2019). Basic principles of the using of medicinal plants and their collections for treatment and prevention of diabetes mellitus type 2. *Phytotherapy chasopys*, 4, 43–46.
- Savych, A., Marchyshyn, S., Harnyk, M., Kudria, V., & Ocheretniuk, A. (2021). Determination of amino acids content in two samples of the plant mixtures by GC-MS. *Pharmacia*, 68(1), 283-289.
- Savych, A., Marchyshyn, S., & Milian, I. (2021). Determination of carbohydrates in the herbal antidiabetic mixtures by GC-MC. *Acta Pharmaceutica*, 71(3), 429-443.
- Savych, A., Marchyshyn, S., Basaraba, R., & Kryskiw, L. (2021). Determination of carboxylic acids content in the herbal mixtures by HPLC. *ScienceRise: Pharmaceutical Science*, 2(30), 33-39.
- Savych, A., Basaraba, R., Muzyka, N., & Ilashchuk, P. (2021). Analysis of fatty acid composition content in the plant components of antidiabetic herbal mixture by GC-MS. *Pharmacia*, 68(2), 433-439.
- Savych, A., Marchyshyn, S., Kozyr, H., & Yarema, N. (2021). Determination of inulin in the

- herbal mixtures by GC-MS method. *Pharmacia*, 68(1), 181-187.
19. Savych, A., Marchyshyn, S., & Basaraba, R. (2020). Determination of fatty acid composition content in the herbal antidiabetic collections. *Pharmacia*, 67(3), 153-159.
 20. Savych, A., Marchyshyn, S., & Doroshenko, O. (2020). Screening of hypoglycemic activity of herbal mixtures (Message 2). *Ukrainian biopharmaceutical journal*, 3(64), 22-29.
 21. Savych, A. O., Marchyshyn, S. M., & Milian, I. I. (2020). Screening study of hypoglycemic activity of herbal mixtures (Presentation 3). *Clinical Pharmacy*, 24(4), 38-46.
 22. Savych, A. O., Marchyshyn, S. M., & Basaraba, R. Yu. (2020). Determination of hypoglycemic activity of the herbal mixtures in screening study. *Pharmacology and Drug Toxicology*, 14(5), 344-351.
 23. WHO Guidelines on good agricultural and mixture practices (GACP) for medicinal plants (2003). *World Health Organization*, Geneva, Switzerland, 72.
 24. Singh, J. (2015). International conference on harmonization of technical requirements for registration of pharmaceuticals for human use. *Journal of pharmacology & pharmacotherapeutics*, 6(3), 185-187.
 25. Ellulu, M. S., Rahmat, A., Patimah, I., Khaza'ai, H., & Abed, Y. (2015). Effect of vitamin C on inflammation and metabolic markers in hypertensive and/or diabetic obese adults: a randomized controlled trial. *Drug design, development and therapy*, 9, 3405-3412.
 26. Govema, P., Bains, G., Borgonetti, V., Cettolin, G., Giachetti, D., Magnano, A. R., et al. (2018). Phytotherapy in the management of diabetes: a review. *Molecules*, 23(1), 105.
 27. May J. M. (2016). Ascorbic acid repletion: A possible therapy for diabetic macular edema? *Free radical biology & medicine*, 94, 47-54.
 28. Cazeau, R. M., Huang, H., Bauer, J. A., & Hoffman, R. P. (2016). Effect of Vitamins C and E on Endothelial Function in Type 1 Diabetes Mellitus. *Journal of diabetes research*, 2016, 3271293.
 29. Shi, L., Du, X., Guo, P., Huang, L., Qi, P., & Gong, Q. (2020). Ascorbic acid supplementation in type 2 diabetes mellitus: A protocol for systematic review and meta-analysis. *Medicine*, 99(45), e23125.




Figure. 1. Absorption maximum of ascorbic acid content in the herbal mixture (Values are expressed as mean \pm SD (n=4))




Figure 2. Calibration curve of standard solution of ascorbic acid

Table 1. Linearity, LOD and LOQ of the spectrophotometric method for the determination of ascorbic acid content

Standard	Range, mg/mL	Regression curve	R ²	LOD, mg/mL	LOQ, mg/ML
L-ascorbic acid	0.02-0.1	y=0.0371x-0.0081	0.9929	0.01	0.04

Note: Results are based on two replicate samples of each standard concentration, each measured twice (n = 4).

Table 2. Precision and accuracy of the spectrophotometric method for the determination of ascorbic acid content

Standard	Concentration, mg/mL	Intra-assay precision*, %	Intermediate precision**, %	Recovery, %
L-ascorbic acid	0.25	1.7	1.8	97.2
	0.50	3.2	4.5	104.2
	1.00	2.4	2.9	105.1

Note:

* intra-assay precision – one analyst in one laboratory, with each concentration made as two replicates and each measured three times (n = 6);

** intermediate precision – two analysts on different days in the same laboratory, with each concentration made as two replicates and each measured three times (n = 6).