

Chlorophyta Spirogyra Ethanolic Extract to the Antifungal Activita of Yeasts and Dermatophytes

**Raghunath D POKHARKAR, *Prasad E FUNDE, Shirish S PINGALE
Sunil L GAVSHETE, Vidhate D KISHOR.**

Department of Chemistry, Arts, Commerce & Science College, Narayangaon, Pune,
Maharashtra India

P.G.S.P. Charitable Foundation Funde Takali Ahmednagar Maharashtra India.

Ravikiran and Tribal Woman Development Institution Maharashtra India

Summary

The Antifungal activity of the ethanol extract of *Chlorophyta Spirogyra* was evaluated at two different concentrations by the diffusion method. The ethanol extract of the *Chlorophyta Spirogyra* shows antifungal activity at varied levels in Yeasts (*Candida albicans*, *C. parapsilosis*) and Dermatophytes (*Trichophyton rubru*, *T. mentagrophytes*). The yeast *Candida albicans* was found to be more active and *C. parapsilosis* was found to be less active in inhibition zone. **Dermatophytes** *Trichophyton rubrum* and *T. mentagrophytes* also shows Antifungal activity. *T. mentagrophytes* was found to be more active than reference inhibition zone.

Keywords *Chlorophyta Spirogyra* , Antifungal activity, extract.

Introduction

Marine medicine represents one of the most important fields of marine medicine all over the world. To promote the proper use of marine medicine and to determine their potential as sources for new drugs, it is essential to study medicinal bacteria. Over the past 20 years, there has been an increased interest in the investigation of natural materials as sources of new antifungal agents. Different extracts from marine medicinal bacteria have been tested to identify the source of the therapeutic effects. As a result some natural products have been approved as new antifungal drugs, but there is still an urgent need to identify novel substances that are active towards pathogens with high resistance. Recently, multiple drug resistance has developed due to indiscriminate use of commercial antifungal drugs commonly used in the treatment of fungal infectious diseases making it a global growing-problem. Isolation of antifungal agents less susceptible to regular antifungal and recovery of increasing resistant isolates during antifungal therapy is rising throughout the world which highlights the need for new principles. Natural products, a new source of antifungal agents shows possibly novel mechanisms of action.

Contrary to the synthetic drugs, antifungal of marine bacteria origin are not associated with many side effects and have an enormous therapeutic potential to heal many infectious diseases. For example, vincristine (an antitumor drug), digitalis (a heart regulator), and ephedrine (a bronchodilator used to decrease respiratory congestion) were all originally discovered through research on marine bacteria. This paper reports to study the antifungal activity of Indian marine bacteria, *Chlorophyta Spirogyra* against an array of human pathogens. The ethanol extract of the *Chlorophyta Spirogyra* was evaluated for the potential antifungal property. The selection of this plant for evaluation was based on its traditional usage. The marine bacteria possesses high content of active compound by HPLC methods and the Antifungal activity of the crude ethanol extract of *Chlorophyta Spirogyra* was evaluated at two different concentrations by the diffusion method.

Materials and Methods

Marine bacteria

Chlorophyta Spirogyra marine bacteria were collected in marine water from Konkan, Maharashtra, India. The dried marine bacteria were homogenized to fine powder and further subjected to extraction.

Crude Extraction


The crude ethanol extract was obtained by extracting 15gm of dried powder in 150 ml ethanol and was kept on a rotary shaker for 24hrs. The extract was filtered, centrifuged at 5000rpm for 15 min and was dried under reduced pressure. The yield obtained for ethanol extract was 40.10% with respect to the initial dry material. The extract was stored at 5°C in airtight bottles.

HPLC spectrum of ethanol extract of *Chlorophyta Spirogyra*

HPLC was applied for testing the presence of number of organic compounds available of ethanol extract of *Chlorophyta Spirogyra*. One of the major organic component with 51.670% and 7.053 retention time may have antifungal activity.

Method

Column : YMC ODS 150 mm × 4.6 mm × 5 μ , ID: E-AC-1/06/COL/013
Mobile Phase:
A: 0.05 % TFA in Water /B: 0.05 % TFA in Acetonitrile
Inj. Vol: 10 μL , Col. Temp.: 30°C, Flow rate: 1.4 mL/min
Gradient:
5% to 95 %B in 8 min, Hold for 1.5 min ,9.51-12 min 05 %B



Antifungal investigation

Antifungal activity of *Chlorophyta Spirogyra* extract was tested by the agar diffusion method described in *European Pharmacopoeia*. Minimum inhibitory concentration (MIC) was determined using the serial two-fold broth dilution method described by Pepeljnjak *et al.*. For the agar diffusion method, 25 ± 2 mL of sterile and melted SGA at $45-50^\circ\text{C}$ was inoculated with 1 mL of approximately $1\text{ to }5 \times 10^6$ CFU mL^{-1} of yeast cells or conidia and mycelial structures of dermatophytes in sterile physiological saline in Petri dishes (9cm). Inoculum density was measured with McFarland's standard solution of freshly prepared barium sulfate in sterile water; density of 0.01% BaCl_2 in 1% H_2SO_4 solution equals approximately 3×10^8 cells mL^{-1} . After drying SGA at room temperature for a maximum of 30 minutes, holes of 6 mm in diameter were made with stainless steel cylinders and filled (60 μL) with the ethanol extract. One experiment was done in a Petri dish for one fungal strain. Plates were then incubated at $+4^\circ\text{C}$ for 1 hour and at $25 \pm 2^\circ\text{C}$ for 48hrs for yeasts and 7 days for dermatophytes. After the incubation period, inhibition zones were measured and expressed in mm. Minimum inhibitory concentration (MIC) and minimum fungicidal concentration (MFC) were determined using the serial broth dilution method described by Pepeljnjak *et al.*. MIC was determined by the broth two-fold macro dilution method in Sabouraud 2% glucose broth starting with 50% (V/V) of the fluid extract or essential oil. MIC is defined as the lowest concentration of extract or essential oil that allows no more than 20% growth of the fungi, visualized as a reduced

number of colonies after removing the loop with approx. 10 µL of each dilution on SGA and incubation at 25 ±2°C for 72 hrs for yeasts and 7 days for dermatophytes. *MFC* is defined as the lowest concentration of the extract that completely inhibited the growth of fungi.

Results and Conclusions

In the *Chlorophyta Spirogyra* sample studied, HPLC was applied for testing the presence of number of organic compounds available of ethanol extract *Chlorophyta Spirogyra* . Antifungal activity is shown in Tables I and II. It shows antifungal activity against all the species of dermatophytes investigated, with inhibition zones from 16 to 17 mm. The largest inhibition zone was observed against *T. mentagrophytes* (25 mm). The ethanol extract *Chlorophyta Spirogyra* of extract had a well-balanced antifungal effect on the fungi species *Candida albicans*, *C. parapsilosis* with an inhibition zone of 16 to 17 mm. The ethanol extract *Chlorophyta Spirogyra* of extract shows antifungal activity against all species of dermatophytes studied, with inhibition zones of 17 mm *Trichophyton rubrum* and 25 mm (*Trichophyton mentagrophytes*), All the fungi investigated are sensitive to the ethanol extract *Chlorophyta Spirogyra* . inhibition of the growth of fungi was observed both on the surface and in the depth of the agar during the experiment when the ethanol extract *Chlorophyta Spirogyra* was dropped into the hole (60 µL) of the Petri dish: concentration of the ethanol extract *Chlorophyta Spirogyra* aerosol in the Petri dish was 0.9 µL cm⁻³, with an incubation temperature of 25±2°C. Data on minimum inhibitory concentrations and minimum fungicidal concentrations for the ethanol extract *Chlorophyta Spirogyra*.

The antifungal activity of the ethanol extract *Chlorophyta Spirogyra* was testes invitro. Yeasts *Candida albicans*, *C. parapsilosis* and Dermatocytes *Trichophyton rubrum*, *T. mentagrophytes* showed more potent antifungal activity. . The extract inhibited the growth of Yeasts and dermatophytes in lower concentrations than the extract itself.

Table I. Antifungal activity of anise ethanol extract *Chlorophyta Spirogyra* by the diffusion method

	Fungi	Inhibition Zone (mm)*	
		Reference substance	ethanol extract
Yeasts	<i>Candida albicans</i>	18 mm	17 mm
	<i>C. parapsilosis</i>	18 mm	16 mm
Dermatophytes	<i>Trichophyton rubrum</i>	21 mm	17 mm
	<i>T. mentagrophytes</i>	24 mm	25 mm

a: Nistatin as reference substance (0.5 mg mL⁻¹) for yeasts and ketoconazole for dermatophytes. b: GPA – growth promotion activity. c: AE – fungicidal influence of aerosol (0.7 µL cm⁻³).

Table II. Antifungal activity of anise ethanol extract *Chlorophyta Spirogyra* by the dilution method

	Fungi	Inhibition Zone (mm)*			
		Reference substance ($\mu\text{g mL}^{-1}$)		ethanol extract (%, v/v)	
		<i>MIC</i> ^b	<i>MFC</i> ^c	<i>MIC</i>	<i>MFC</i>
Yeasts	<i>Candida albicans</i>	0.5	1	18	23
	<i>C. parapsilosis</i>	1	2	17	24
Dermatophytes	<i>Trichophyton rubrum</i>	0.25	0.5	1.4	3
	<i>T. mentagrophytes</i>	0.50	1	8.0	17

a Nistatin as reference antimycotic against yeasts tested and ketoconazole against dermatophytes tested.

b *MIC* – minimum inhibitory concentration.

c *MFC* – minimum fungicidal concentration.

d Significantly different from fluid extract ($p < 0.01$).

nt – not tested.

References

1. Shahidi, G.H.; Karimi Nik, A. (2004). Antibacterial activity of some medicinal plants of Iran against *Pseudomonas aeruginosa* and *P. fluorescens*. *Asian J. Plant Sci.*, 3, 61-64.
2. Tortora, G.J.; Funke, B.R.; Case, C.L. (2001). Microbiology: an introduction. Benjamin Cummings, San Francisco, p.88.
3. Vlietinck, A.J.; Van Hoof, L.; Totte, J.; Lasure, A.; Vanden Berghe, D.; Rwangobo, P.C.; Mvukiyuniwami, J. (1995). Screening of hundred Rwandese medicinal plants for antimicrobial and antiviral properties. *J. Ethnopharmacol.*, 46, 31-47.
4. Yoshida, T.; Chou, T.; Haba, K.; Okano, Y.; Shingu, T.; Miyamoto, K.; Koshiura, R.; Okuda, T. (1989). Camellin B and nobotanin I, macrocyclic ellagitannin dimmers and related dimmers, and their antitumor activity. *Chem. Pharm. Bull.*, 37, 3174-3176
5. Barbour, E.K.; Al Sharif, M.; Sagherian, V.K.; Habre, A.N.; Talhouk, R.S.; Talhouk, S.N.; (2004). Screening of selected indigenous plants of Lebanon for antimicrobial activity. *J. Ethnopharmacol.*, 93, 1-7.

6. Cho, T.; Koshiur, R.; Miyamoto, K.; Nitt, A.; Okud, T.; Yoshida, T.; (1990). Woodfordin C, a macro-ring hydrolyzable tannin dimer with antitumor activity, and accompanying dimers from *Woodfordia fruticosa* flowers. *Chem. Pharm. Bull.*, 38, 1211-1217.
7. Cragg, G.M.; Newman, D.J.; Snader, K.M.; (1997). Natural products in drug discovery and development. *J. Nat. Prod.*, 60, 52-60.
8. El-Faky, F.K.; Attif, O.; Aboul Ela, M.; Gaanem, N.; (1995). Antimicrobial evaluation of extracts from some Yemeni plants. *Alexandrian J. Pharm. Sci.*, 9, 35-37.
9. Farnsworth, N.R.; Morris, R.W.; (1976). Higher plants: the sleeping giant of drug development. *Am. J. Pharm.*, 48, 46-52.
10. Hamil, F.A.; Apio, S.; Mubiru, N.K.; Bukenya-Ziraba, R.; Mosango, M.; Maganyi, O.W.; Soejarto, D.D. (2003). Traditional herbal drugs of Southern Uganda, II: literature analysis and antimicrobial assays. *J. Ethnopharmacol.*, 84, 57-78.
11. Machado, T.B.; Pinto, A.V.; Pinto, M.C.F.R.; Leal, I.C.R.; Silva, M.G.; Amaral, A.C.F.; Kuster, R.M.; Netto-dosSantos, K.R.; (2003). *In vitro* activity of Brazilian medicinal plants, naturally occurring naphthoquinones and their analogues, against methicillin-resistant *Staphylococcus aureus*. *Int. J. Antimicrob.*, 21, 279-284.
12. Motsei, M.L.; Lindsey, K.L.; van Staden, J.; Jaeger, A.K.; (2003). Screening of traditionally used South African plants for antifungal activity against *Candida albicans*. *J. Ethnopharmacol.*, 86, 235- 241.
13. Al-Dubai, A.S.; Al-Khulaidi, A.A.; (1996). Medical and aromatic plants of Yemen (in Arabic). Obadi Center for Studies and Publishing, Sana'a, Yemen.
14. Anjaria, J.; Parabia, M.; Dwivedi, S.; (2002). Ethnovet heritage Indian ethnoveterinary medicine – An Overview,; Pathik Enterprise, Ahmedabad (Gujarat), India.