

**Antimicrobial and Antifungal Activities of Local Edible Fern
Stenochlaena Palustris (Burm. F.) Bedd**

Z. Zuraini¹, S. Sasidharan², S. Roopin Kaur¹ and M. Nithiyayini¹

¹School of Distance Education, Universiti Sains Malaysia, 11800 USM, Pulau Pinang, Malaysia.

²Institute for Research in Molecular Medicine (INFORM), Universiti Sains Malaysia, 11800 USM, Malaysia.

*Corresponding author: Phone +60125323462, email: srisasidharan@yahoo.com

Summary

This current research was conducted to study the uninvestigated (or less investigated) medicinal ferns found in Malaysia in the search for more effective and non-toxic pharmacological natural compounds. Methanol of extracts from winged bean *Stenochlaena palustris* root, stem and leaf extracts were tested for their antimicrobial activity against 15 microbial species, including 10 bacterial pathogens, one yeasts, and four molds using the disk diffusion assay technique. The leaf extract was found to be most effective against all of the tested organisms, followed by the stem and root extracts. The minimum inhibitory concentrations (MICs) of the leaf extracts determined by the broth dilution method ranged from 50 to 12.5 mg/mL. The preliminary results of present investigation appear to indicate that *S. palustris* of Malaysian Edible fern have higher potential antimicrobial properties.

Key words: antimicrobial activity, antifungal activity, fern, *Stenochlaena palustris*

Introduction

Infectious diseases and their control become serious problem in the medical field. Antibiotics usually suggested for the treatment of infectious diseases, have never been pleasing because of their toxic effect and exert a negative impact on the consumer. As an alternative strategy to prevent infectious diseases, natural compound of plants are being tested for their antimicrobial activity and serve as template for new and more effective antimicrobial agents. This research attempted to study uninvestigated (or less investigated) medicinal plants and ferns found in Malaysia in the search for potential, more effective and non-toxic pharmacological natural compounds.

The pteridophytes which constitute ferns and ferns allies, have been known to man for more than 2000 years, and also been mentioned in ancient literature (1, 2). It has been observed that pteridophytes are not infected by microbial pathogens, which may be one of the important factors for the evolutionary success of pteridophytes and the fact that they survived for more than 350 million years.

This information's encouraged us to further investigate other tropical fern found in Malaysia for antimicrobial activity. *Stenochlaena palustris* (Burm.) Bedd. (Pteridaceae) is a fern trailing over the ground or scrambling high up trees. It is endemic to a large part of tropical areas from southern and northern India through Malaysia to Polynesia and Australia (3). The tender leaves of *S. palustris* are used as a contraceptive by the local people in the central district province of Papua New Guinea (PNG) and in the Nicobar Islands (4, 5). A search for alkaloid-containing plants in New Guinea found the leaves of *S. palustris* to be alkaloid-negative (6). No other chemical studies on this species have been reported. Therefore, the current study was conducted to study the antimicrobial activity of methanolic extract of *S. palustris* against bacterial and fungal strains.

Materials and methods

Materials: The root, stem and leaf of *S. palustris* were collected from Seberang Jaya, Penang, Malaysia, in Mac 2009.

Extraction of plant material: Fern sample was extracted by cold percolation in methanol (consecutively three times) at room temperature for 24 h the resultant extracts were filtered and concentrated to dryness under reduced pressure below 40 °C in rotary evaporator.

Antimicrobial activity: Antibacterial and antifungal activities of the fern extracts were investigated by the disk diffusion method (7, 8). The MHA plates, containing an inoculum size of 10^6 colony-forming units (CFU)/mL of bacteria or 2×10^5 CFU/mL yeast cells or molds spores on SDA and PDA plates, respectively, were spread on the solid plates with an L-shaped glass rod. Then disks (6.0-mm diam.) impregnated with 25 μ L of each extract at a concentration of 100.0 mg/mL were placed on the inoculated plates. Similarly, each plate carried a blank disk by adding every solvent control alone in the center, and antibiotic disks (6.0-mm diam.) of 30 μ g/mL Chloramphenicol (for bacteria), and 30 μ g/mL Miconazole nitrate (for fungi) were also used as positive controls. All of the plates were incubated at 37°C for 18 hours for bacteria and at 28°C for 48 hours for fungi. The zones of growth inhibition around the disks were measured after 18 hours of in incubation at 37°C for bacteria and 48 hours for fungi at 28°C, respectively.

MIC was determined by both broth dilution methods. For broth dilution tests, 0.1mL of standardized suspension of bacteria (10^6 CFU/mL) and fungal cell or spores (5×10^5 CFU/mL) was added to each tube (containing fractions of three extracts at a final concentration of 0 to 20.0mg/mL) and incubated at 37°C for bacteria for 18 hours or at 28°C for fungi for 48 hours. MICs were taken as the average of the lowest concentration showing no growth of the organism and the highest concentration showing visible growth by macroscopic evaluation. Each assay was performed in triplicate.

Results and discussion

The antimicrobial activity data of methanol extract of *S. palustris* are shown in Tables 1 and 2. All the extracts were found to be active in at least one of the microbial strains tested. In general, both bacteria and fungi were found to be susceptible to the test agents than fungi. The preliminary disk diffusion assay (Table 1) of *S. palustris* extracts of root, stem and leaf showed that the leaf extract had the most distinct effect on most of the tested microorganisms, followed by the stem and root extracts. However, two commercial antibiotics were more effective than any of the extract tested. The minimum inhibitory concentration (MIC) values of methanolic extract of *S. palustris* leaf was evaluated and summarized in Table 2. The determination of MIC of root extract was carried out in this study since the extract showed the best activity in the preliminary antimicrobial screening assay. The methanolic extracts of *S. palustris* root gave low MIC values against *Enterobacter aerogenes*, *Salmonella typhi* and *Azospirillum lipoferum* with concentration of 12.5 mg/ml. However, the *Klebsiella pneumoniae* showed a highest MIC value of 50.0 mg/ml when tested with root extract of *S. palustris*. The MIC values of the extracts seem to be relatively higher. However, being crude extracts, the overall antimicrobial activity screening results are only indicative of the potential of this fern extracts for the medicinal purposes.

The preliminary results of present investigation appear to indicate that *S. palustris* of Malaysian Edible fern especially lower groups (pteridophytes) have higher potential antimicrobial properties. Further studies aimed at the isolation and identification of active substances from the methanol extracts of *S. palustris* may disclose other compounds with better value for antimicrobial agents.

References

1. Kirtikar KR, Basu BD. Indian Medicinal Plants, L.M. Basu, Allahabad, 1935, p. 2752.
2. Nayar BK. Medicinal Ferns of India, *Bulletin of Natural Botanic Gardens*. 1957; 29: 1-36.
3. Holttum RE. A Revised Flora of Malaya; Government Printing Office, Singapore, 1968, Vol 2: p 412.
4. Wolff-Eggert R. Über Heilpflanzen von Papua-Neuguinea. Ph.D. Thesis, Friedrich-Alexander-Universität, Erlangen-Nürnberg, 1977; p 142.
5. Dagar H S. *J. Econ. TaxBot.* 1989; 13: 395–397.
6. Hertley TG, Dunstone EA, Fitzgerald JS, Johns SR, Lamberton JA. *J. Nat. Prod.* 1973; 36: 217–319.
7. Bauer AW, Kirby WMM, Sherris JC, and Turck M. Antibiotic susceptibility testing by standardized single disc method. *Am. J. Clin. Pathol.* 1966; 36: 493-496.
8. Alzoreky NS and Nakahara K. Antibacterial activity of extracts from some edible plants commonly consumed in Asia. *Int. J. Food Microbiol.* 2003; 80:223-230.

Table 1: Antimicrobial activity (zone of inhibition)^a of *Stenochlaena palustris*

Microorganisms	Zone of inhibition (mm) ^b				
	Leaves	Stems	Roots	C	M
BACTERIA					
Gram-positive					
<i>Staphylococcus aureus</i>	14	13	15	24	ND
<i>Bacillus subtilis</i>	10	9	18	22	ND
<i>Micrococcus sp</i>	9	16	8	26	ND
Gram-negative					
<i>Enterobacter aerogenes</i>	16	10	13	25	ND
<i>E. coli</i>	18	14	15	22	ND
<i>Proteus mirabilis</i>	17	24	21	22	ND
<i>Klebsiella pneumoniae</i>	16	17	14	25	ND
<i>Salmonella typhi</i>	–	9	9	21	ND
<i>Azospirillum lipoferum</i>	14	18	20	26	ND
<i>Azobacter</i>	18	12	17	23	ND
FUNGUS					
<i>Penizillium chrysogenum</i>	14	13	9	ND	21
<i>Rhizopus stolonifer</i>	–	–	–	ND	20
<i>Aspergillus niger</i>	11	15	16	ND	24
<i>Fusarium sp.</i>	12	8	10	ND	22
<i>Saccharomyces cerevisiae</i>	15	11	16	ND	21

^aDisc diffusion technique

^bThe values (average of triplicate) are diameters of zone of inhibition at 100mg/mL of crude extract, 30µg/mL of Chloramphenicol and 30µg/mL of Miconazole nitrate

ND : Not Determined

C : Chloramphenicol

M : Miconazole nitrate

Table 2: Determination of MIC values of root extracts of *Stenochlaena palustris*

Bacteria	Concentration (mg/ml)											
	100.000	50.000	25.000	12.500	6.250	3.125	1.563	0.781	0.391	0.195	0.098	
Gram-positive												
<i>Staphylococcus aureus</i>	-	-	-	+	+	+	+	+	+	+	+	+
<i>Bacillus subtilis</i>	-	-	-	+	+	+	+	+	+	+	+	+
<i>Micrococcus sp</i>	-	-	-	+	+	+	+	+	+	+	+	+
Gram-negative												
<i>Enterobacter aerogenes</i>	-	-	-	-	+	+	+	+	+	+	+	+
<i>E. coli</i>	-	-	-	-	+	+	+	+	+	+	+	+
<i>Proteus mirabilis</i>	-	-	-	+	+	+	+	+	+	+	+	+
<i>Klebsiella pneumoniae</i>	-	-	+	+	+	+	+	+	+	+	+	+
<i>Salmonella typhi</i>	-	-	-	-	+	+	+	+	+	+	+	+
<i>Azospirillum lipoferum</i>	-	-	-	-	+	+	+	+	+	+	+	+
<i>Azobacter</i>	-	-	-	+	+	+	+	+	+	+	+	+

- : Indicates absence of growth

+ : Indicates presence of growth