

**THIN LAYER CHROMATOGRAPHIC PROFILING AND EVALUATION OF  
ANALGESIC ACTIVITY OF *MUSA PARADISIACA* LEAF EXTRACTS IN MICE**

**Aniket Tarafdar<sup>1</sup>, Sanjib Bhattacharya<sup>2\*</sup>, J. N. Pandey<sup>1</sup>, Moulisha Biswas<sup>1</sup>**

<sup>1</sup>*Bengal Institute of Pharmaceutical Sciences, Kalyani, Nadia 741235, West Bengal, India*

<sup>2</sup>*Bengal School of Technology, Delhi Road, Sugandha, Hooghly 712102, West Bengal, India*

**\*For correspondence:** E-mail: sakkwai@yahoo.com

**Summary**

*Musa paradisiaca* (L.) (Musaceae) commonly known as banana in English, is a perennial tree like herb grown indigenously in tropics and subtropics and also cultivated commercially in India for its fruits. The present study assessed the different solvent extracts of *M. paradisiaca* leaf for thin layer chromatography (TLC) and also evaluated their analgesic potential by acetic acid induced writhing assay in Swiss albino mice. The chloroform extract yielded maximum spots in TLC, followed by petroleum ether and methanol extracts respectively. All of the test extracts exhibited significant analgesic activity. The methanol extract was found to be the most potent followed by the chloroform and petroleum ether extracts respectively. The present preliminary study demonstrates marked analgesic activity of *M. paradisiaca* leaf.

**Key words:** Analgesic, *Musa paradisiaca*, writhing, leaf.

**Introduction**

Traditional medicine worldwide is being re-evaluated by extensive research on different plant species and their therapeutic principles. The rich wealth of plant kingdom can represent a novel source of new compounds with therapeutic activity. *Musa paradisiaca* (L.) (Musaceae) commonly known as banana in English or *Kola-gachh* in Bengali and *Kela* in Hindi, is a perennial tree like herb grown indigenously in tropics and subtropics. It is widely found throughout Indian subcontinent and commercially cultivated in various parts of India for its consumable fruits. All parts of this plant have been traditionally used in India for several medicinal purposes. Ayurvedic physicians of Karnataka and Kerala states of India recommended *M. paradisiaca* for the treatment of urinary stones (1). The stem juices of *M. paradisiaca* have been reported for dissolving pre-formed stones and in preventing the

formation of stones in the urinary bladder of rats (2, 3). Its stem juice has also been used in nervous affectations like epilepsy, hysteria and in dysentery and diarrhoea. Several oligosaccharides comprising fructose, xylose, galactose, glucose and mannose occur naturally in banana, making it an excellent prebiotic for the selective growth of beneficial bacteria in the intestine (4). Previous workers reported several experimental studies on *Musa paradisiaca*. Among them the potential use of banana fruit pulp to treat ulcers have been explored by a number of investigators (5-8). However, its chromatographic profile and analgesic assessment are still not reported. Therefore, in the present investigation we attempted these studies on the leaf extracts of *M. paradisiaca*.

### **Materials and methods**

**Plant material:** The mature leaves of *Musa paradisiaca* (L.) (Musaceae) were collected during November 2010 from Nadia, West Bengal, India. The plant material was taxonomically identified at the Central National Herbarium, Botanical Survey of India, Howrah, West Bengal, India. The voucher specimen [CNH/I-I/(83)/2010/Tech.II/352] was maintained in our research laboratory for future reference. The plant material was shade-dried with occasional shifting and then powdered with mechanical grinder, passing through sieve no. 40, and stored in an air-tight container.

**Preparation of plant extracts:** The dried powdered material (350 g) was defatted with petroleum ether (60-80 °C), the percentage extractive value was 3.72 % w/w. The defatted powder material thus obtained was further extracted with chloroform and methanol for 72 h in a percolator. The solvent was distilled off in reduced pressure and resulting semisolid mass was vacuum dried using rotary flash evaporator to yield a solid residue and the percentage extractive values were accordingly 5.09 % w/w and 7.87 % w/w respectively. The preliminary phytochemical analysis was performed for all three extracts to identify the phytoconstituents present in the extracts (9).

**Chemicals:** Acetyl salicylic acid (aspirin) and glacial acetic acid from Sigma Chemical Co. Ltd. (St. Louis, MO, USA). All other chemicals and reagents were of analytical grade obtained commercially.

#### **Experimental animals**

Adult male albino mice of Swiss strain weighing  $20 \pm 2$  g were procured from registered breeders (Rita Ghosh & Co., Kolkata, India) and maintained under standard laboratory conditions (temperature  $25 \pm 2^\circ\text{C}$  with dark and light cycle 14/10 h). They were allowed free access to standard dry pellet diet (Hindustan Lever, Kolkata, India) and water *ad libitum*. The mice were acclimatized to laboratory condition for 10 days before commencement of the experiment.

**Thin layer chromatographic studies:** Each solvent extract was subjected to thin layer chromatography (TLC) as per standard one dimensional ascending method. The results and chromatograms are depicted in Figures 1-3.

#### **Analgesic evaluation: acetic acid-induced writhing test**

Swiss albino mice were divided into five groups ( $n = 6$ ). Group I received acetic acid (1% v/v, 10 ml/kg b.w., i.p.) and writhing reflex was noted for the period of 15 minutes. Group II received aspirin (100 mg/kg b.w. p.o.) Group III, IV and V received the petroleum ether, chloroform and methanol extracts at the doses of 200 mg/kg b.w., p.o. respectively. 30 min

after aspirin and extracts administration, group II to V received acetic acid (1% v/v, 10 ml/kg b.w., i.p.) and writhing reflex was noted for the period of 15 min (10). The mean writhing scores in each group were calculated and expressed the percentage of protection using the following formula:-

$$(\text{Control mean} - \text{Treated mean} / \text{Control mean}) \times 100 \%$$

**Table 1.** Analgesic effect of extracts from *M. paradisiaca* on acetic acid induced writhing in mice.

Treatment	Dose	Number of writhes	% Protection
Acetic acid (1% v/v)	10 ml/kg	52.83 ±1.400	-
Acetic acid + Aspirin	100 mg/kg	17.66 ±1.606*	66.57
Acetic acid + Pet. ether extract of <i>M. paradisiaca</i>	200 mg/kg	25.09 ±1.261*	52.50
Acetic acid + Chloroform extract of <i>M. paradisiaca</i>	200 mg/kg	20.05±1.371*	62.04
Acetic acid + Methanol extract of <i>M. paradisiaca</i>	200 mg/kg	18.89±1.471*	64.24

Values are mean ± SEM ( $n = 6$ ). \*  $p < 0.001$  when compared to control.

## Results and discussion


Preliminary phytochemical studies showed the presence of steroids in the petroleum ether extract; steroids, triterpenoids, saponins, glycosides and carbohydrates in the chloroform extract; and steroids, saponins, glycosides and carbohydrates in the methanol extract.

Among the various methods for separating plant constituents, the thin layer chromatographic procedure is the one of the most commonly used techniques of general application (11). Thin layer chromatography (TLC) involves the separation of mixtures of organic compounds on thin layers of adsorbents that are usually coated on glass, plastic, or aluminium sheets; and this particular technique is the easiest, cheapest and most widely used method for the characterization of natural products and their preparations (12). The chloroform extract yielded maximum spots in TLC, followed by petroleum ether and methanol extracts respectively (Figures 1-3). All of these TLC profiles may serve as characteristic fingerprint of *M. paradisiaca* leaf. These data would therefore be suitable for monitoring the identity and purity of the plant material and for detecting adulterations and substitutions.


The analgesic efficacy of *M. paradisiaca* leaf extracts was evaluated by acetic acid induced writhing method in mice to assess peripheral (non-narcotic) type of analgesic activity (13). Acetic acid induced writhing is chemically induced nociception by intraperitoneal injection of dilute acetic acid solution to mice. The chemical agents can produce nociceptive reactions in mice. Intra-peritoneal injection of phenyl para quinone, bradykinin or dilute acetic acid (1-3% v/v) produces pain reaction that is characterized as writhing response. Constriction of abdomen, turning of trunk (twist) and extension of hind limbs (at least one) are considered as writhing reaction to chemically induced pain (13, 14).

Acetic acid induced writhing test is known as a visceral pain model nociception. Several mediators like kinins, acetylcholine, substance P, calcitonin-gene-related peptide and prostaglandins (PG) take part in visceral pain model nociception and transmission of the nociception from the viscera. In this test both central and peripheral analgesics are detected. Analgesics of both narcotic (central) e. g. morphine, pentazocin, pethidine and non-narcotic (peripheral) type, e. g. aspirin, ibuprofen, indomethacin can inhibit the writhing response in mice (13-15).


Based on the data obtained from the present study, all the extracts had effective peripheral analgesic actions. The methanol extract was found to be the most potent followed by the chloroform and petroleum ether extracts respectively (Table 1). The present preliminary study confirms marked analgesic activity of *M. paradisiaca* leaf which may be due to presence of multitude of constituents as revealed by TLC profile.



**Fig. 1.** TLC profile of the pet. ether extract of *M. paradisiaca* leaf. Solvent system: benzene: chloroform: ethyl acetate (5: 3: 2).  $R_f$  values: 0.40, 0.60, 0.64, 0.74, 0.84, 0.92.



**Fig. 2.** TLC profile of the chloroform extract of *M. paradisiaca* leaf. Solvent system: benzene: chloroform: ethyl acetate (2: 3: 5).  $R_f$  values: 0.12, 0.16, 0.28, 0.41, 0.52, 0.62, 0.74, 0.78, 0.84.



**Fig. 3.** TLC profile of the methanol extract of *M. paradisiaca* leaf. Solvent system: benzene: chloroform: ethyl acetate (5: 3: 2).  $R_f$  values: 0.15, 0.30, 0.55, 0.87, 0.96.

### Acknowledgement

The authors are thankful to Mr. Subir K. Pal, President & CEO of Bengal Institute of Pharmaceutical Sciences for his kind co-operation and financial support to complete this work successfully.

**References**

1. Pellai N, Aashan TN. Ayurveda Prakashika. S.T. Reddiar and Son, Vidyarambham Press, Quilon, 1955, 97.
2. Kailash. P, Bharathi K, Srinivasan. K. Evaluation of Musa (Paradisiaca Linn. Cultivar) "Puttubale" stem juice for antilithiatic activity in albino rats. *Ind J Physiol and Pharmacol* 1993; 37: 337-41.
3. Kailash P, Varalakshmi P. Effect of banana stem juice on biochemical change in liver of normal and hyperoxaluric rats. *Ind J Exp Biol* 1992; 30: 440-2.
4. Gibson N. Dietary modulation of the human gut micro flora using prebiotics. *Britain J Nutrition* 1998; 80(4)S: 209-12.
5. Elliot RC, Heward E.JF. The influence of a banana supplemented diet on gastric ulcers in mice. *Pharmacol Research Communication* 1976; 8:167-71.
6. Best R. The antiulcerogenic activity of the unripe plantain banana. *Brit J Pharmacol* 1984; 82: 107-16.
7. Goel S. Stimulation of gastric and colonic mucosal eicosonoid synthesis by plantain banana. *J Pharmceutial and Pharmacol* 1989; 41: 747-50.
8. Lewis G, Shaw H. A natural flavonoid and synthetic analogues protect the gastric mucosa from aspirin-induced erosion. *J Nutrition Biochem* 2001; 12(2): 95-100.
9. Kokate CK (1994): *Practical Pharmacognosy*. 4th Edition. New Delhi. Vallabh Prakashan. pp. 107-112.
10. Kostar R., Anderson M., de Beer E.J. Acetic acid for analgesic screening. *Fed. Proc.* 1959; 18: 412.
11. Kokate CK, Purohit AP, Gokhale SB. *Pharmacognosy*. 34th ed. Nirali Prakashan, Pune, 2006, 114.
12. Stahl E. Apparatus and General Techniques. In: Stahl E, editor. *Thin Layer Chromatography, A Laboratory Handbook*. 2nd ed. Springer-Verlag, Berlin, Heidelberg, New York, 1969, 52-85.
13. Vogel H.G. (ed.). *Drug Discovery and Evaluation, Pharmacological Assays*. 2<sup>nd</sup> ed.: Springer Verlag: Berlin, Heidelberg, 2002.
14. Kulkarni S.K. *Hand Book of Experimental Pharmacology*. 3rd ed. Vallabh Prakashan: New Delhi, 1999.
15. Parkes, M.W., J.T. Pickens. 1965. Conditions influencing the inhibition of analgesic drugs of the response to intra peritoneal injections of phenylbenzoquine in mice. *Brit. J. Pharmacol.*, 25: 81-87.